



12.16.99

A/ box seg

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.

BB1205 US NA

First Named Inventor or Application Identifier

SAVERIO CARL FALCO

Express Mail Label No.

EK369475295US

Express Mailing Date

DECEMBER 15, 1999

12/15/99

09/464528

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents

Box Patent Application

Washington, DC 20231

☒

Fee (Authority to charge deposit account below.)

(Submit an original, and a duplicate for fee processing)

☒

Specification

(preferred arrangement set forth below)

[Total Pages

27]

- Descriptive title of the invention

- Cross References to Related Applications (if needed)

- Statement Regarding Fed sponsored R & D (if needed)

- Reference to Microfiche Appendix (if filed)

- Background of the Invention

- Brief Summary of the Invention

- Brief Description of the Drawings (if filed)

- Detailed Description

- Claim(s)

- Abstract of the Disclosure

6.

☐ Microfiche Computer Program (Appendix)

7.

Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)

a.

☒ Computer Readable Copy

b.

☒ Paper Copy (identical to computer copy)

c.

☒ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8.

☐ Power of Attorney

9.

☐ Information Disclosure
Statement (IDS)/Cover
Letter plus PTO-1449☐Copies of IDS
Citations

10.

☐ Preliminary Amendment

11.

☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)

12.

☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)

13.

☒ Other:GENE SEQUENCE LISTING - 12 PAGES

3.

☒

Drawing(s) (35 USC 113)

[Total Sheets

10]

4.

☐

Oath or Declaration

[Total Pages

0]

a.

☐ Newly executed (original or copy)

b.

☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 14 completed)

i.

☐ DELETION OF INVENTORS
Signed Statement below at 15 deleting
inventor(s) named in the prior application,
see 37 CFR 1.63(d)(2) and 1.33(b).

5.

☐Incorporation by Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a
copy of the oath or declaration is supplied under Box 4b, is
considered as being part of the disclosure of the
accompanying application and is hereby incorporated by
reference therein.

14. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐

Continuation

☐

Divisional

☐

Continuation-in-part (CIP)

of prior Application No.: ____/____

15.

☐**DELETION OF INVENTOR(S) STATEMENT:** This application is being filed by less than all the inventors named in the prior application. In accordance with 37 CFR 1.63(d)(2) and 1.33(b), the Assistant Commissioner is requested to delete the name(s) of the following person or persons who are not inventors of the invention being claimed in this application:

16.

☒

Amend the specification by inserting before the first line the sentence:

-- This application claims benefit priority of Provisional Application No. 60/113,045, filed December 21, 1998. --

17.

☐

Cancel in this application original claims ____ of the prior application before calculating the filing. (At least one original independent claim must be retained for filing purposes.)

18.

☐

Priority of foreign Application No. _____ filed on _____ in

_____ is claimed under 35 U.S.C. 119.
(country)

CLAIMS	(1) FOR	(2) NUMBER FILED	(3) NUMBER EXTRA	(4) RATE	(5) CALCULATIONS
	TOTAL CLAIMS (37 CFR 1.16(c))	28 - 20 =	8	x \$ 18 =	\$ 144.00
	INDEPENDENT CLAIMS (37 CFR 1.16(b))	2 - 3 =	0	x \$ 78 =	0
	MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$ 260 =	260.00
				BASIC FEE (37 CFR 1.16(a))	+ \$ 760.00
				TOTAL =	\$ 1164.00

19. The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Account No. 04-1928:

a. ☒ Fees required under 37 CFR 1.16.

b. ☒ Fees required under 37 CFR 1.17.

20. ☐ Other:

21. CORRESPONDENCE ADDRESS

NAME	Lynne M. Christenbury				
ADDRESS	E. I. du Pont de Nemours and Company				
	Legal - Patents 1007 Market Street				
CITY	Wilmington	STATE	Delaware	ZIP CODE	19898
COUNTRY	U.S.A.	TELEPHONE	302-992-5481	FAX	302-892-7949

22. SIGNATURE OF ATTORNEY OR AGENT REQUIRED

NAME	Lynne M. Christenbury	REG. NO.: 30,971
SIGNATURE	<i>Lynne M. Christenbury</i>	
DATE	15 December 1999	

TITLES-ADENOSYL-L-METHIONINE SYNTHETASE PROMOTER
AND ITS USE IN EXPRESSION OF TRANSGENIC GENES IN PLANTSFIELD OF THE INVENTION

5 This invention relates to a plant promoter, in particular, to an S-adenosyl-L-methionine synthetase (SAMS) promoter and subfragments thereof and their use in regulating the expression of at least one heterologous nucleic acid fragment in plants.

BACKGROUND OF THE INVENTION

Recent advances in plant genetic engineering have opened new doors to engineer
10 plants having improved characteristics or traits, such as, resistance to plant diseases, insect resistance, herbicidal resistance, enhanced stability or shelf-life of the ultimate consumer product obtained from the plants and improvement of the nutritional quality of the edible portions of the plant. Thus, a desired gene (or genes) from a source different than the plant, but engineered to impart different or improved characteristics or qualities, can be
15 incorporated into the plant's genome. This new gene (or genes) can then be expressed in the plant cell to exhibit the new trait or characteristic.

In order to obtain expression of the newly inserted gene in the plant cell, the proper regulatory signals must be present and be in the proper location with respect to the gene. These regulatory signals include a promoter region, a 5' non-translated leader sequence and a
20 3' transcription termination/polyadenylation sequence.

A promoter is a DNA sequence that directs cellular machinery of a plant to produce RNA from the contiguous coding sequence downstream (3') of the promoter. The promoter region influences the rate, developmental stage, and cell type in which the RNA transcript of the gene is made. The RNA transcript is processed to produce messenger RNA (mRNA)
25 which serves as a template for translation of the RNA sequence into the amino acid sequence of the encoded polypeptide. The 5' non-translated leader sequence is a region of the mRNA upstream of the protein coding region that may play a role in initiation and translation of the mRNA. The 3' transcription termination/polyadenylation signal is a non-translated region downstream of the protein coding region that functions in the plant cells to cause termination
30 of the RNA transcript and the addition of polyadenylate nucleotides to the 3' end of the RNA.

It has been shown that certain promoters are able to direct RNA synthesis at a higher rate than others. These are called "strong promoters". Certain other promoters have been shown to direct RNA production at higher levels only in particular types of cells or tissues and are often referred to as "tissue specific promoters". In this group, many seed storage
35 protein genes' promoters have been well characterized and widely used, such as the phaseolin gene promoter of *Phaseolus vulgaris*, the helianthinin gene of sunflower, the β -conglycinin gene of soybean (Chen et al., (1989) *Dev. Genet.* 10, 112-122), the napin gene promoter of *Brassica napus* (Ellerstrom et al, (1996) *Plant Mol. Biol.* 32, 1019-1027), the oleosin gene

promoters of *Brassica* and *Arabidopsis* (Keddie et al, (1994) *Plant Mol. Biol.* 24, 327-340; Li, (1997) Texas A&M Ph.D. dissertation, pp. 107-128; Plant et al, (1994) *Plant Mol. Biol.* 25, 193-205). Another class of tissue specific promoters is described in, U.S. Patent No. 5,589,583, issued to Klee et al. on December 31, 1996; these plant promoters are capable of conferring high levels of transcription of chimeric genes in meristematic tissues and/or rapidly dividing cells. In contrast to tissue-specific promoters, "inducible promoters" direct RNA production in response to certain environmental factors, such as heat shock, light, hormones, ion concentrations etc. (Espartero et al, (1994) *Plant Mol. Biol.* 25, 217-227; Gomez-Gomez and Carrasco, (1998) *Plant Physiol.* 117, 397-405; Holtorf et al, (1995) *Plant Mol. Biol.* 29, 637-646; MacDowell et al, (1996) *Plant Physiol.* 111, 699-711; Mathur et al, (1992) *Biochem. Biophys. Acta* 1137, 338-348; Mett et al, (1996) *Transgenic Res.* 5, 105-113; Schoffl et al, (1989) *Mol. Gen. Genet.* 217, 246-253; Ulmasov et al, (1995) *Plant Physiol.* 108, 919-927).

Promoters that are capable of directing RNA production in many or all tissues of a plant are called "constitutive promoters". The ideal constitutive promoter should be able to drive gene expression in all cells of the organism throughout its development. Expression of many so-called constitutive genes, such as actin (McDowell et al., (1996) *Plant Physiol.* 111, 699-711; Wang et al., (1992) *Mol. Cell Biol.* 12, 3399-3406), and ubiquitin (Callis et al, (1990) *J. Biol. Chem.* 265, 12486-12493; Rollfinke et al, (1998) *Gene* 211, 267-276) varies depending on the tissue types and developmental stages of the plant. The most widely used constitutive promoter, the cauliflower mosaic virus 35S promoter, also shows variations in activity in different plants and in different tissues of the same plant (Atanassova et al., (1998) *Plant Mol. Biol.* 37, 275-285; Battraw and Hall, (1990) *Plant Mol. Biol.* 15, 527-538; Holtorf et al., (1995) *Plant Mol. Biol.* 29, 637-646; Jefferson et al., (1987) *EMBO J.* 6, 3901-3907; Wilmink et al., (1995) *Plant Mol. Biol.* 28, 949-955). The cauliflower mosaic virus 35S promoter is also described in U.S. Patent No. 5,106,739. The tissue-specific expression and synergistic interactions of sub-domains of the promoter of cauliflower mosaic virus are discussed in U.S. Patent No. 5,097,025, which issued to Benfey et al. on March 17, 1992. A *Brassica* promoter (hsp80) that provides for constitutive expression of heterologous genes in a wide range of tissues and organs is discussed in U.S. Patent No. 5,612,472 which issued to Wilson et al. on March 18, 1997.

Since the patterns of expression of a chimeric gene (or genes) introduced into a plant are controlled using promoters, there is an ongoing interest in the isolation and identification of novel promoters which are capable of controlling expression of a chimeric gene or (genes).

SUMMARY OF THE INVENTION

This invention concerns an isolated nucleic acid fragment comprising a promoter wherein said promoter consists essentially of the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15, or 16 or said promoter consists essentially of a fragment or subfragment that is

substantially similar and functionally equivalent to the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15, or 16.

In a second embodiment, this invention concerns a chimeric gene comprising at least one heterologous nucleic acid fragment operably linked to the promoter of the invention.

In a third embodiment, this invention concerns plants containing this chimeric gene and seeds obtained from such plants.

In a fourth embodiment, this invention concerns a method of increasing or decreasing the expression of at least one heterologous nucleic acid fragment in a plant cell which comprises:

- (a) transforming a plant cell with the chimeric gene described above;
- (b) growing fertile mature plants from the transformed plant cell of step (a);
- (c) selecting plants containing the transformed plant cell wherein the expression of the heterologous nucleic acid fragment is increased or decreased.

In a fifth embodiment, this invention concerns an isolated nucleic acid fragment comprising a constitutive plant SAMS promoter.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCES

The invention can be more fully understood from the following detailed description, the drawings and the Sequence Descriptions that form a part of this application. The Sequence Descriptions contain the three letter codes for amino acids as defined in 37 C.F.R. §§ 1.821-1.825, which are incorporated herein by reference.

SEQ ID NO:1 is the nucleotide sequence comprising the entire cDNA insert in clone s2.12b06 which encodes a soybean S-adenosyl-L-methionine synthetase protein.

SEQ ID NO:2 is the nucleotide sequence comprising a soybean S-adenosyl-L-methionine synthetase genomic DNA fragment.

SEQ ID NO:3 is the nucleotide sequence of a portion of the cDNA insert in clone srr1c.pk002.b21 encoding a portion of a soybean S-adenosyl-L-methionine synthetase protein.

SEQ ID NO:4 is a 32 base oligonucleotide primer, designated sam-5, used to amplify the soybean S-adenosyl-L-methionine synthetase promoter region via PCR.

SEQ ID NO:5 is a 24 base oligonucleotide primer, designated sam-6, used to amplify the soybean S-adenosyl-L-methionine synthetase promoter region via PCR.

SEQ ID NO:6 is the nucleotide sequence comprising a soybean S-adenosyl-L-methionine synthetase promoter fragment produced via PCR using primers sam-5 (SEQ ID NO:4) and sam-6 (SEQ ID NO:5).

SEQ ID NO:7 is a 22 base oligonucleotide primer, designated sam-9, used to amplify the soybean S-adenosyl-L-methionine synthetase promoter region via PCR.

SEQ ID NO:8 is a 19 base oligonucleotide primer, designated atps-9, used to amplify a chimeric gene comprising a SAMS promoter fragment and a portion of the ATP sulfurylase (ATPS) gene via PCR.

SEQ ID NO:9 is a 21 base oligonucleotide primer, designated cgs-8, used to amplify a chimeric gene comprising a SAMS promoter and a portion of the cystathionine- γ -synthase 1 (CGS1) gene via PCR.

SEQ ID NO:10 is a 20 base oligonucleotide antisense primer, designated atps-4, used to amplify the ATP sulfurylase transcript via RT-PCR.

SEQ ID NO:11 is a 21 base oligonucleotide antisense primer, designated cgs-10, used to amplify the cystathionine- γ -synthase 1 transcript via RT-PCR.

SEQ ID NO:12 is a 20 base oligonucleotide primer, designated atps-3, used to amplify an ATP sulfurylase cDNA via PCR.

SEQ ID NO:13 is a 23 base oligonucleotide primer, designated cgs-9, used to amplify a cystathionine- γ -synthase 1 cDNA via PCR.

SEQ ID NO:14 is a 2165 nucleotide sequence comprising a soybean S-adenosyl-L-methionine synthetase genomic DNA fragment which starts at the 5' end of SEQ ID NO:2, and ends at the ATG translation start codon of the S-adenosyl-L-methionine synthetase.

SEQ ID NO:15 is a 1574 nucleotide sequence comprising a DNA fragment which starts at the 5' end of SEQ ID NO:2, and ends at the ATG translation start codon of the S-adenosyl-L-methionine synthetase, and wherein a 591 nucleotide intron sequence has been removed.

SEQ ID NO:16 is a 719 nucleotide sequence comprising a DNA fragment which starts at nucleotide 4 of SEQ ID NO:6, and ends at the ATG translation start codon of the S-adenosyl-L-methionine synthetase, and wherein a 591 nucleotide intron sequence has been removed.

SEQ ID NO:17 is a 6975 nucleotide sequence comprising plasmid pMH40 Δ .

SEQ ID NO:18 is a 3985 nucleotide sequence comprising a SAMS promoter::GUS::3' Nos DNA fragment present in plasmid pZSL11.

SEQ ID NO:19 is a 3684 nucleotide sequence comprising a SAMS promoter::ATPS::3' Nos DNA fragment.

SEQ ID NO:20 is a 3963 nucleotide sequence comprising a SAMS promoter::CGS1::3' Nos DNA fragment.

Figures 1A and 1B depict Southern hybridization analyses of SAMS genes. Soybean genomic DNA was digested with BamHI, EcoRI, HindIII, KpnI, and SacI, and then the blot was hybridized with a full length SAMS cDNA (SEQ ID NO:1) probe in Figure 1A or with a SAMS promoter fragment (SEQ ID NO:6) probe in Figure 1B.

Figure 2 depicts a SAMS genomic DNA sequence (SEQ ID NO:2) and the alignment of the overlapping region with SAMS cDNA sequence (SEQ ID NO:1). The 2336 bp SAMS

genomic DNA sequence has a 191 bp region aligned with the 5' end sequence of the SAMS cDNA with six mismatches. The region used to make the SAMS promoter by adding the NcoI site at its 3' end is underlined. The translation start codon is in bold.

Figure 3 depicts the structure of the SAMS::GUS expression cassette. The SAMS promoter was cloned into pMH40Δ to replace its 35S promoter. The structure of the resulted SAMS::GUS construct was generated by Vector NTI™ software (InforMax, Inc., North Bethesda, MD).

Figure 4 depicts a histochemical GUS expression analysis of transgenic *Arabidopsis* plants harboring the SAMS::GUS expression cassette. *Arabidopsis* tissues were incubated at 37°C with X-Gluc overnight and dehydrated with ethanol. (A) Flower buds; (B) leaf; (C) Inflorescence stem and a cauline leaf; (D, E, F) developing siliques; (G) Developing seeds and embryos. All of the seeds were derived from GUS-positive siliques. Genetic segregation of the GUS gene was demonstrated by the blue funiculus of the white seed in the right upper corner.

Figure 5 depicts a fluorometric GUS expression assay of transgenic *Arabidopsis* plants harboring the SAMS::GUS expression cassette. Triple samples of flowers, leaves, stems, siliques coats, young seeds, medium seeds, old seeds, and dry seeds collected from SAMS::GUS transgenic *Arabidopsis* plants were assayed for GUS activity. The graph was generated by Microsoft Excel and the standard deviation is indicated by the upper part of each column.

Figure 6 depicts a histochemical GUS transient expression analysis of SAMS promoter in corn. The pZSL11 (SAMS::GUS) or the pMH40Δ (35S::GUS) plasmid DNA was delivered into corn callus (A, C) or leaf discs (B, D), and the GUS activity was detected by incubation with X-Gluc overnight at 37° C. (A, B) Transformed with pZSL11 DNA; (C, D) Transformed with pMH40Δ DNA.

Figures 7(A) and 7(B) depict the presence and expression of transgenic soybean ATPS and CGS1 genes controlled by the SAMS promoter in transgenic *Arabidopsis* plants. Figure 7(A) is a PCR analysis. Genomic DNA of ten transgenic *Arabidopsis* plants (1 to 10), wild type *Arabidopsis* (a), wild type soybean (s), and plasmid DNA of SAMS::CGS1 or SAMS::ATPS in binary vectors (p) were used as templates in PCR with gene-specific primers. PCR of ten SAMS::CGS1 transgenic plants with primer sam-9 which is specific to SAMS promoter, and primer cgs-8 which is specific to soybean CGS1 (upper). PCR of ten SAMS::ATPS transgenic plants with primer sam-9 which is specific to SAMS promoter, and primer atps-1 which is specific to soybean ATPS gene (lower). Figure 7(B) is an RT-PCR analysis. Total leaf RNA of ten transgenic *Arabidopsis* plants (1 to 10), wild type *Arabidopsis* (a), and wild type soybean (s) were used as templates in RT-PCR with gene-specific primers. First strand cDNA was synthesized from a gene-specific antisense primer with reverse transcriptase, and then the first strand cDNA was amplified by PCR with both

sense and antisense primers. RT-PCR of ten SAMS::CGS1 transgenic plants with primers, cgs-9 (sense) and cgs-10 (antisense), specific to soybean CGS1 gene (upper). RT-PCR of ten SAMS::ATPS transgenic plants with primers, atps-3 (sense) and atps-4 (antisense), specific to soybean ATPS gene (lower).

Figure 8 depicts induction of SAMS promoter activity by methionine. Seeds of ten transgenic Arabidopsis lines transformed with SAMS::GUS construct were germinated on filter papers soaked with H₂O, 1x Murashige and Skoog salt, 0.01 mM, and 1 mM methionine. Ten days old seedlings were harvested and assayed for GUS activity. The solid bar and hollow bar indicate, respectively, the average and the standard variation of three samples for each treatment.

Figure 9 depicts a northern hybridization. Soybean total RNAs from leaves, roots, stems, young seeds, medium seeds, old seeds, and pod coats (L, R, S, Y, M, O, and P) were used to make the RNA blot which was hybridized with a full length SAMS cDNA (SEQ ID NO:1) probe.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized.

As used herein, an “isolated nucleic acid fragment” is a polymer of ribonucleotides (RNA) or deoxyribonucleotides (DNA) that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA.

The terms “subfragment that is functionally equivalent” and “functionally equivalent subfragment” are used interchangeably herein. These terms refer to a portion or subsequence of an isolated nucleic acid fragment in which the ability to alter gene expression or produce a certain phenotype is retained whether or not the fragment or subfragment encodes an active enzyme. For example, the fragment or subfragment can be used in the design of chimeric genes to produce the desired phenotype in a transformed plant. Chimeric genes can be designed for use in co-suppression or antisense by linking a nucleic acid fragment or subfragment thereof, whether or not it encodes an active enzyme, in the appropriate orientation relative to a plant promoter sequence.

The terms “substantially similar” and “corresponding substantially” as used herein refer to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate gene expression or produce a certain phenotype. These terms also refer to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially alter the functional properties of the resulting nucleic acid fragment relative to the initial, unmodified fragment. It is therefore understood, as those skilled in the art will appreciate, that the invention encompasses more than the specific exemplary sequences.

Moreover, the skilled artisan recognizes that substantially similar nucleic acid sequences encompassed by this invention are also defined by their ability to hybridize, under moderately stringent conditions (for example, 0.5 X SSC, 0.1% SDS, 60° C) with the sequences exemplified herein, or to any portion of the nucleotide sequences reported herein and which are functionally equivalent to the promoter of the invention. Preferred substantially similar nucleic acid sequences encompassed by this invention are those sequences that are 80% identical to the nucleic acid fragments reported herein or which are 80% identical to any portion of the nucleotide sequences reported herein. More preferred are nucleic acid fragments which are 90% identical to the nucleic acid sequences reported herein, or which are 90% identical to any portion of the nucleotide sequences reported herein. Most preferred are nucleic acid fragments which are 95% identical to the nucleic acid sequences reported herein, or which are 95% identical to any portion of the nucleotide sequences reported herein. Sequence alignments and percent similarity calculations may be determined using the Megalign program of the LASARGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences are performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments and calculation of percent identity of protein sequences using the Clustal method are KTUPLE=1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. For nucleic acids these parameters are GAP PENALTY=10, GAP LENGTH PENALTY=10, KTUPLE=2, GAP PENALTY=5, WINDOW=4 and DIAGONALS SAVED=4. A "substantial portion" of an amino acid or nucleotide sequence comprises enough of the amino acid sequence of a polypeptide or the nucleotide sequence of a gene to afford putative identification of that polypeptide or gene, either by manual evaluation of the sequence by one skilled in the art, or by computer-automated sequence comparison and identification using algorithms such as BLAST (Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410) and Gapped Blast (Altschul, S. F. et al., (1997) *Nucleic Acids Res.* 25:3389-3402); see also www.ncbi.nlm.nih.gov/BLAST/).

"Gene" refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. "Native gene" refers to a gene as found in nature with its own regulatory sequences. "Chimeric gene" refers to any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature. Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature. "Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that

is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

A “heterologous nucleic acid fragment” refers to a nucleic acid fragment comprising a nucleic acid sequence that is different from the nucleic acid sequence comprising the plant promoter of the invention.

“Coding sequence” refers to a DNA sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include, but are not limited to, promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989, *Biochemistry of Plants* 15:1-82). It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of some variation may have identical promoter activity. An “intron” is an intervening sequence in a gene that is transcribed into RNA but is then excised in the process of generating the mature mRNA. The term is also used for the excised RNA sequences. An “exon” is a portion of the sequence of a gene that is transcribed and is found in the mature messenger RNA derived from the gene, but is not necessarily a part of the sequence that encodes the final gene product.

The “translation leader sequence” refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA

stability or translation efficiency. Examples of translation leader sequences have been described (Turner, R. and Foster, G. D. (1995) *Molecular Biotechnology* 3:225).

The “3' non-coding sequences” refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., (1989) *Plant Cell* 1:671-680.

“RNA transcript” refers to a product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When an RNA transcript is a perfect complementary copy of a DNA sequence, it is referred to as a primary transcript or it may be a RNA sequence derived from posttranscriptional processing of a primary transcript and is referred to as a mature RNA. “Messenger RNA” (“mRNA”) refers to RNA that is without introns and that can be translated into protein by the cell. “cDNA” refers to a DNA that is complementary to and synthesized from an mRNA template using the enzyme reverse transcriptase. The cDNA can be single-stranded or converted into the double-stranded by using the klenow fragment of DNA polymerase I. “Sense” RNA refers to RNA transcript that includes mRNA and so can be translated into protein within a cell or *in vitro*. “Antisense RNA” refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks expression or transcripts accumulation of a target gene (U.S. Patent No. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, *i.e.* at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. “Functional RNA” refers to antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

“Sense” RNA refers to RNA transcript that includes the mRNA and so can be translated into protein by the cell. “Antisense RNA” refers to a RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. Patent No. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, *i.e.*, at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. “Functional RNA” refers to antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term “operably linked” refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (*i.e.*, that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term “expression”, as used herein, refers to the production of a functional end-product. Expression or overexpression of a gene involves transcription of the gene and translation of the mRNA into a precursor or mature protein. “Antisense inhibition” refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. “Overexpression” refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. “Co-suppression” refers to the production of sense RNA transcripts capable of suppressing the expression or transcript accumulation of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020). The mechanism of co-suppression may be at the DNA level (such as DNA methylation), at the transcriptional level, or at post-transcriptional level.

“Altered expression” refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ significantly from the amount of the gene product(s) produced by the corresponding wild-type organisms.

“Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. The preferred method of corn cell transformation is use of particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook, J., Fritsch, E. F. and Maniatis, T., *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1989 (hereinafter “Sambrook et al., 1989”) or Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. and Struhl, K. (eds.), *Current Protocols in Molecular Biology*, John Wiley and Sons, New York, 1990 (hereinafter “Ausubel et al., 1990”).

“PCR” or “Polymerase Chain Reaction” is a technique for the synthesis of large quantities of specific DNA segments, consists of a series of repetitive cycles (Perkin Elmer Cetus Instruments, Norwalk, CT). Typically, the double stranded DNA is heat denatured, the two primers complementary to the 3' boundaries of the target segment are annealed at low temperature and then extended at an intermediate temperature. One set of these three consecutive steps comprises a cycle.

An “expression construct” is a plasmid vector or a subfragment thereof comprising the instant chimeric gene. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and

patterns of expression (Jones et al., (1985) *EMBO J.* 4:2411-2418; De Almeida et al., (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

Although the SAMS enzyme is present in most plant cell types, no SAMS promoter capable of driving gene expression in most or all plant cell types has been described. Previous studies indicated that plants contain multiple SAMS genes which are differentially expressed in response to various stresses (Schroder *et al.* (1997) *Plant Mol. Biol.* 33:211-222). A SAMS promoter that is preferentially active in a particular tissue type, i.e. vascular (Peleman *et al.*, (1989) *Plant Cell* 1, 81-93; Mijnsbrugge *et al.*, (1996) *Plant Cell Physiol.* 37, 1108-1115), was also known. However, it was not possible to predict, before the studies reported herein, whether any SAMS gene was controlled by a constitutive promoter. It is demonstrated herein that constitutive SAMS promoters do, in fact, exist in plants, and that such promoters can be readily isolated and used by one skilled in the art.

This invention concerns an isolated nucleic acid fragment comprising a constitutive plant SAMS promoter. This invention also concerns an isolated nucleic acid fragment comprising a promoter wherein said promoter consists essentially of the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15 or 16 or said promoter consists essentially of a fragment or subfragment that is substantially similar and functionally equivalent to the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15 or 16. A nucleic acid fragment that is functionally equivalent to the instant SAMS promoter is any nucleic acid fragment that is capable of controlling the expression of a coding sequence or functional RNA in a similar manner to the SAMS promoter. The expression patterns of the SAMS promoter are defined in the following paragraphs.

Northern-blot hybridization experiments indicated that SAMS gene transcripts are present in a variety of soybean tissues and that the abundance of SAMS gene transcripts does not differ greatly from tissue to tissue (Figure 9 and Example 3). Strong expression of the SAMS gene was also inferred by the high frequency of occurrences of cDNA sequences with homology to SAMS (ESTs) in a soybean cDNA sequence database created by sequencing random cDNAs from libraries prepared from many different soybean tissues. ESTs encoding SAMS can be easily identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database, e.g., SAMS from *Oryza sativa* (EMBL Accession No. Z26867) or SEQ ID NO:1 provided herein. SAMS homologs were among the most abundant classes of cDNAs found in the soybean libraries. This indicated that SAMS was a highly expressed

gene in most soybean cell types. The data obtained from sequencing many SAMS ESTs also indicated that there were several SAMS isoforms encoded by the soybean genome.

A soybean cDNA clone designated s2.12b06 was found to encode a protein which is very similar to the protein encoded by the cDNA to *Oryza sativa* SAMS (pLog value for this match was 61.59). The soybean cDNA clone designated s2.12b06 was completely sequenced (SEQ ID NO:1) and found to contain an opening reading frame which encodes a full length SAMS polypeptide. Southern hybridization analysis of soybean genomic DNA with this full length SAMS cDNA as a probe suggested that there are approximately four related SAMS genes in the soybean genome (Figure 1A), which is consistent with the EST sequencing data.

The soybean SAMS cDNA clone was used to isolate a soybean genomic DNA fragment containing more than 2000 nucleotides upstream (5') of the SAMS protein coding sequence by hybridization of a soybean genomic DNA library to the SAMS cDNA fragment probe. Southern hybridization analysis of soybean genomic DNA using a 1314 base pair DNA fragment from upstream of the SAMS protein coding sequence as a probe indicated that this fragment is unique in the soybean genome (Figure 1B).

The promoter activity of the soybean genomic DNA fragment upstream of the SAMS protein coding sequence was assessed by linking the fragment to a reporter gene, the *E. coli* β -glucuronidase gene (GUS) (Jefferson (1987) Plant Mol. Biol. Rep. 5:387-405), transforming the SAMS promoter::GUS expression cassette into *Arabidopsis*, and analyzing GUS expression in various cell types of the transgenic plants. GUS expression was detected in all parts of the transgenic plants that were analyzed. These results indicated that the nucleic acid fragment contained a constitutive promoter. Since SAMS catalyzes the reaction to synthesize S-adenosyl-L-methionine from methionine and ATP, free methionine levels might regulate SAMS promoter activity. To see if the SAMS promoter is regulated by external methionine, the SAMS::GUS transgenic *Arabidopsis* seeds were germinated in the presence or absence of methionine. Ten day old seedlings were analyzed for GUS activity according to the protocol described in Example 5. Ten independent transgenic lines were tested and all of them responded similarly. GUS activity was more than two-fold higher in seedlings germinated in the presence of methionine (Figure 8). The increased SAMS promoter activity in the presence of methionine may be particularly useful for efforts to increase methionine biosynthesis via overexpression of enzymes in the methionine biosynthetic pathway or the sulfate assimilation pathway. It is clear from the disclosure set forth herein that one of ordinary skill in the art could readily isolate a constitutive plant SAMS promoter from any plant by performing the following procedure:

1) obtaining a SAMS cDNA from a desired plant by any of a variety of methods well known to those skilled in the art including, but not limited to, (a) random sequencing of ESTs from a cDNA library and characterizing the ESTs via a BLAST search as described

above; or (b) hybridizing a cDNA library to a known plant SAMS cDNA; or (c) PCR amplification using oligonucleotide primers designed from known SAMS cDNAs;

2) obtaining a genomic DNA fragment that includes approximately 500 to 3000 nucleotides from the region 5' to a SAMS protein coding sequence, which contains a SAMS promoter, by hybridization of a genomic DNA library to a SAMS cDNA fragment probe;

3) operably linking the nucleic acid fragment containing the region upstream (5') of the SAMS protein coding sequence to a suitable reporter gene ; there are a variety of reporter genes that are well known to those skilled in the art, including the bacterial GUS gene, the firefly luciferase gene, and the green fluorescent protein gene; any gene for which an easy and reliable assay is available can serve as the reporter gene

4) transforming a chimeric SAMS promoter::reporter gene expression cassette into an appropriate plant for expression of the promoter. There are a variety of appropriate plants which can be used as a host for transformation that are well known to those skilled in the art, including the dicots, *Arabidopsis*, tobacco, soybean, oilseed rape, peanut, sunflower, safflower, cotton, tomato, potato, cocoa and the monocots, corn, wheat , rice, barley and palm. The terms "oilseed rape" and "oilseed Brassica" are used interchangeably herein.

5) testing for expression of a SAMS promoter in various cell types of transgenic plants, e.g., leaves, roots, flowers, seeds, transformed with the chimeric SAMS promoter::reporter gene expression cassette by assaying for expression of the reporter gene product. A constitutive SAMS promoter will produce high level expression of the reporter in all, or nearly all, of the plant tissues tested.

In another aspect, this invention concerns a chimeric gene comprising at least one heterologous nucleic acid fragment operably linked to the promoter of the present invention. Chimeric genes can be constructed by operably linking the nucleic acid fragment of the invention, i.e., the SAMS promoter or a fragment or a subfragment that is substantially similar and functionally equivalent to any portion of the nucleotide sequence set forth in SEQ ID NOS:6, 14, 15 or 16, to a heterologous nucleic acid fragment. Any heterologous nucleic acid fragment can be used to practice the invention. The selection will depend upon the desired application or phenotype to be achieved. The various nucleic acid sequences can be manipulated so as to provide for the nucleic acid sequences in the proper orientation.

Plasmid vectors comprising the instant chimeric genes can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host cells. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene.

The plasmid vectors or chimeric genes can be used to transform plant cells. Transformation techniques are well known to those skilled in art as discussed above. A preferred method of plant cell transformation is the use of particle-accelerated or "gene gun"

transformation technology (Klein *et al.* (1978) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050). The chimeric gene will normally be joined to a marker for selection in plant cells. The marker may be resistance to a biocide, particularly an antibiotic, such as kanamycin, G418, bleomycin, hygromycin, chloramphenicol, or the like. The particular marker employed will be one which will allow for selection of transformed cells as compared to cells lacking the heterologous nucleic acid sequence which has been introduced. Examples of plant cells which can be transformed using plant transformation techniques include, but are not limited to, monocot and dicot plant cells such as soybean, oilseed *Brassica* species, corn, peanut, rice, wheat, sunflower, safflower, cotton, cocoa, tobacco, tomato, potato, barley, palm, Arabidopsis and the like.

In addition to the bacterial GUS gene, two soybean genes, ATP sulfurylase (ATPS) and cystathionine- β -synthase 1 (CGS1), were also successfully expressed by this promoter in transgenic Arabidopsis, as depicted in Figure 7. This further validates the application of the SAMS promoter of the invention in plant genetic engineering practice.

The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression of the chimeric genes (Jones *et al.*, (1985) *EMBO J.* 4:2411-2418; De Almeida *et al.*, (1989) *Mol. Gen. Genetics* 218:78-86). Thus, multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by northern analysis of mRNA expression, western analysis of protein expression, or phenotypic analysis. Also of interest are seeds obtained from transformed plants displaying the desired expression profile.

The level of activity of the SAMS promoter is comparable to that of many known strong promoters, such as the CaMV 35S promoter (Atanassova *et al.*, (1998) *Plant Mol. Biol.* 37:275-285; Battraw and Hall, (1990) *Plant Mol. Biol.* 15:527-538; Holtorf *et al.*, (1995) *Plant Mol. Biol.* 29:637-646; Jefferson *et al.*, (1987) *EMBO J.* 6:3901-3907; Wilmink *et al.*, (1995) *Plant Mol. Biol.* 28:949-955), the *Arabidopsis* oleosin promoters (Plant *et al.*, (1994) *Plant Mol. Biol.* 25:193-205; Li, (1997) Texas A&M University Ph.D. dissertation, pp. 107-128), the *Arabidopsis* ubiquitin extension protein promoters (Callis *et al.*, 1990), a tomato ubiquitin gene promoter (Rollfinke *et al.*, 1998), a soybean heat shock protein promoter (Schoffl *et al.*, 1989), and a maize H3 histone gene promoter (Atanassova *et al.*, 1998).

Expression of the chimeric genes in most plant cell makes the SAMS promoter of the instant invention especially useful when constitutive expression of a target heterologous nucleic acid fragment is required. Examples of suitable target heterologous nucleic acid fragments include, but are not limited to, a herbicide-resistance or pathogen-resistance nucleic acid fragment. Another useful feature of the constitutive plant SAMS promoter is its expression profile in developing seeds. The SAMS promoter of the invention is most active in developing seeds at early stages and gradually turns down at later stages. Such activity is

indicated by the GUS activity detected in seeds of transgenic *Arabidopsis* plants containing a SAMS::GUS expression cassette as shown in Figures 4 and 5. The expression profile of the claimed SAMS promoter is different from that of many seed-specific promoters, e.g., seed storage protein promoters, which often provide highest activity in later stages of development (Chen *et al.*, (1989) *Dev. Genet.* 10:112-122; Ellerstrom *et al.*, (1996) *Plant Mol. Biol.* 32:1019-1027; Keddle *et al.*, (1994) *Plant Mol. Biol.* 24:327-340; Plant *et al.*, (1994) *Plant Mol. Biol.* 25:193-205; Li, (1997) Texas A&M University Ph.D. dissertation, pp. 107-128). Thus, the SAMS promoter will be a very attractive candidate when overexpression of a gene in embryos is desired at an early developing stage. For example, it may be desirable to overexpress a gene regulating early embryo development or a gene involved in the metabolism prior to seed maturation.

One general application of the SAMS promoter of the invention is to construct chimeric genes that can be used in the selection of transgenic cell lines in plant transformation. Currently, many of the selectable marker genes for plant transformation are under the control of the cauliflower mosaic virus 35S promoter. Since the SAMS promoter of the invention is active in seedlings and callus, the appropriate selection phase for transgenic plants or cell lines, this promoter may be used as an alternative to the 35S promoter to drive the expression of selectable marker genes.

Another general application of the SAMS promoter of the invention is to construct chimeric genes that can be used to reduce expression of at least one heterologous nucleic acid fragment in a plant cell. To accomplish this a chimeric gene designed for cosuppression of a heterologous nucleic acid fragment can be constructed by linking the fragment to the SAMS promoter of the present invention. (See U.S. Patent No. 5,231,020 for methodology to block plant gene expression via cosuppression.) Alternatively, a chimeric gene designed to express antisense RNA for a heterologous nucleic acid fragment can be constructed by linking the fragment in reverse orientation to the SAMS promoter of the present invention. (See U.S. Patent No. 5,107,065 for methodology to block plant gene expression via antisense RNA.) Either the cosuppression or antisense chimeric gene can be introduced into plants via transformation. Transformants wherein expression of the heterologous nucleic acid fragment is decreased or eliminated are then selected.

This invention also concerns a method of increasing or decreasing the expression of at least one heterologous nucleic acid fragment in a plant cell which comprises:

- (a) transforming a plant cell with the chimeric genes described herein;
- (b) growing fertile mature plants from the transformed plant cell of step (a);
- (c) selecting plants containing a transformed plant cell wherein the expression of the heterologous nucleic acid fragment is increased or decreased.

Transformation and selection can be accomplished using methods well-known to those skilled in the art including, but not limited to, the methods described herein.

EXAMPLES

The present invention is further defined in the following Examples. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

Unless otherwise stated, all parts and percentages are by weight and degrees are Celsius. Techniques in molecular biology were typically performed as described in Ausubel, F. M., et al., (1990, Current Protocols in Molecular Biology, John Wiley and Sons, New York) or Sambrook, J. et al., (1989, Molecular cloning - A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

EXAMPLE 1

Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from soybean tissues were prepared in Uni-ZAP XR™ vectors according to the manufacturer's protocol (Stratagene, La Jolla, CA). Conversion of the Uni-ZAP XR™ libraries into plasmid libraries was accomplished according to the protocol provided by Stratagene. Upon conversion, cDNA inserts were contained in the plasmid vector pBluescript™ (Stratagene). DNA was prepared for sequencing from randomly selected bacterial colonies containing recombinant pBluescript™ plasmids either by amplifying the cDNA inserts via polymerase chain reaction using primers specific for vector sequences flanking the cloning site or by preparing plasmid DNA from cultured bacterial cells. Amplified insert DNAs or plasmid DNAs were sequenced in dye-primer sequencing reactions using a Perkin Elmer Model 377 fluorescent sequencer to generate partial cDNA sequences termed expressed sequence tags or "ESTs" (see Adams, M. D. et al., (1991) *Science* 252:1651).

EXAMPLE 2

Identification of SAMS cDNA Clones

ESTs encoding SAMS were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul, S. F., et al., (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish, W. and States, D. J. (1993) *Nature Genetics* 3:266-272 and Altschul, S. F., et al. (1997) *Nucleic Acids Res.*

25:3389-3402) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as “pLog” values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST “hit” represent homologous proteins.

The BLASTX search using the nucleotide sequence from clone s2.12b06 revealed that this nucleotide sequence encoded a protein that was similar to the protein encoded by the cDNA to *Oryza sativa* (EMBL Accession No. Z26867) S-adenosylmethionine synthetase; the pLog value for this match was 61.59. This cDNA clone was completely sequenced (SEQ ID NO:1) and found to contain an opening reading frame ranging from nucleotides 74 to 1252 which is predicted to encode a full length SAMS polypeptide.

A high level of expression of the SAMS genes was inferred by the high frequency of occurrences of soybean cDNA sequences with homology to *Oryza sativa* SAMS obtained from many different cDNA libraries prepared from many different soybean cell types. SAMS homologs were the third most abundant class of ESTs found in the soybean libraries. Although the ranking might not represent a precise estimate of the relative abundance of the SAMS transcripts *in vivo* in all soybean libraries, due to the selective use of different cDNA libraries, it did indicate that SAMS was a highly expressed gene. The EST sequence data also revealed that there were several SAMS isoforms in the soybean genome.

EXAMPLE 3

S-adenosylmethionine Synthetase is Encoded by a Gene Family

Southern hybridization analysis of soybean genomic DNA with a full length SAMS cDNA (SEQ ID NO:1) as a probe suggested that there are at least four related SAMS genes in the soybean genome (Figure 1A). The DNA probe for Southern hybridization was prepared as follows: plasmid DNA was prepared from an overnight bacteria culture in LB broth (GIBCO BRL, Gaithersburg, MD) using QIAprep™ miniprep kit (Qiagen, Valencia, CA); cDNA inserts encoding SAMS were excised by restriction enzyme digestion and recovered from agarose gel following electrophoretic separation using QIAquick™ gel extraction kit (Qiagen). The 1518 bp SAMS cDNA fragment (SEQ ID NO:1) was labeled with digoxigenin-dUTP as a probe by random primed DNA labeling (Boehringer Mannheim). Twenty micrograms of soybean genomic DNA was digested with different restriction enzymes and the resulted fragments were resolved on a 0.7% agarose gel. The DNA gel was depurinated in 0.25 M HCl, denatured in 0.5 M NaOH/1.5 M NaCl, neutralized in 1 M Tris-Cl, pH 8.0/1.5 M NaCl, and transferred in 20x SSC (GIBCO BRL) to nylon membrane (Boehringer Mannheim). The Southern blot was hybridized with the SAMS cDNA-specific probe at 45°C overnight in Easy Hyb (Boehringer Mannheim). The blot was washed 10 minutes in 2xSSC/0.1% SDS, and 3x 10 minutes in 0.1x SSC/0.1% SDS at 65°C. The

hybridized probe was detected with chemiluminescent reagent CDP-Star (Boehringer Mannheim) according to the manufacturer's protocol. Multiple bands were detected in BamHI, EcoRI, and HindIII digestions (Figure 1A). The large band in KpnI and SacI digestions may represent more than one DNA fragment because the band is too big for good resolution. The hybridization patterns presented in Figure 1A and the analysis of partial SAMS cDNA sequences from DuPont's EST database suggest that there are at least four copies of the SAMS gene in the soybean genome and that their sequences are conserved.

The 1314 bp SAMS promoter fragment (SEQ ID NO:6) was labeled with digoxigenin-dUTP also by random primed DNA labeling (Boehringer Mannheim). The labeled SAMS promoter probe was used to hybridize the same Southern blot as above described. The SAMS promoter-specific probe hybridized to a single band in each of the five different digestions, BamHI, EcoRI, HindIII, KpnI, and SacI (Figure 1B). The results indicate that the SAMS promoter has only a single copy in soybean genome.

A northern hybridization experiment indicated that SAMS gene transcripts were present in a variety of soybean tissues and that the abundance of SAMS gene transcripts did not differ greatly from tissue to tissue. Total RNAs were extracted from soybean leaves, stems, young seeds, medium seeds, old seeds, and pod coats using Trizol™ Reagent according to the manufacturer's protocol (GIBCO BRL). Ten micrograms of total RNA were loaded in each well of a 1.2% agarose gel containing 7% formaldehyde in 1x MOPS buffer, 20 mM 3-[N-morpholino]propane-sulfonic acid, 5 mM sodium acetate, 1 mM EDTA, pH 6.0. RNA was transferred to nylon filters (Micron Separations Inc., Westborough, MA) in 10X SSC and crosslinked to the filters with UV light. Filters were hybridized with probes prepared from cDNA insert fragments in 50% deionized formamide, 5x SSPE, 1x Denhardt's solution, 0.1% SDS, and 100 µg denatured salmon sperm DNA (Sigma, St. Louis, MO) at 42° for 24 hours. Filters were washed in 2x SSPE and 0.1% SDS at room temperature for 10 minutes, 1x SSPE and 0.1% SDS at 65° for 10 minutes, and then in 0.1x SSPE and 0.1% SDS at 65° for 10 minutes. Filters were exposed to Kodak X-ray film at -80. The abundance of SAMS transcripts in leaves, roots, stems, young seeds, medium seeds, old seeds, and pod coats can be seen in Figure 9. The weak signals observed in the hybridizations to RNA samples from root and young seed were attributed to underloading, because hybridizations with ribosomal RNAs that serve as internal controls were also relatively weak in those samples (data not shown). Because of the high sequence similarities among the four SAMS gene isoforms, this RNA gel blot was not able to indicate how the isoforms were distributed in any particular tissue. However, the experiment demonstrated that all examined soybean tissues contained SAMS messenger RNA.

EXAMPLE 4

Cloning of the Soybean S-adenosylmethionine Synthetase Gene Promoter

The soybean full length SAMS cDNA (SEQ ID NO:1), obtained in Example 2, was used to generate a probe to isolate a SAMS promoter. The full length SAMS cDNA sequence consisted of 1518 bp, and it had a 74 bp 5'-untranslated region and a PstI site at position 296. Because the cDNA clone was harbored in a pBluescript™ SK vector having a PstI site upstream of the EcoRI cloning site, digestion of the clone with PstI generated a 315 bp fragment of DNA. The resulting restriction fragment contained 19 bp of vector and cloning linker adapter sequence in addition to the 296 bp of SAMS cDNA sequence. This PstI fragment was labeled with α -³²P-dCTP, as described in Example 3, and used as a probe to screen a soybean genomic DNA library that had been constructed in a EMBL3 SP6/T7 vector (ClonTech, Palo Alto, CA). The library was plated with LE392 (ClonTech) cells at 50,000 plaque forming units (pfu) per 150 mm NZCYM agar plate (GIBCO BRL). Plaques were transferred to Hybond nylon membranes, and the plaque replicas were then denatured and neutralized according to the manufacturer (Amersham Life Science, Arlington Heights, IL). The phage DNA was fixed on the membranes by UV-crosslinking (Stratagene). After prehybridization at 65° for 1 hour in 0.5 M NaHPO₄, pH 7.2, 1 mM EDTA, 1% crystalline BSA (Sigma), and 7% SDS, the SAMS 315 bp PstI fragment probe was denatured in boiling water bath for 5 minutes and added to the same hybridization solution, and was hybridized at 65° for 24 hours. The membranes were washed in 40 mM NaHPO₄, pH 7.2, 1 mM EDTA, 0.5% crystalline BSA, and 5% SDS for 10 minutes at room temperature, and then 3x 10 minutes at 65° in 40 mM NaHPO₄, pH 7.2, 1 mM EDTA, and 1% SDS. The membranes were exposed to Kodak X-ray film (Sigma) at -80°. Positive SAMS genomic DNA phage clones were suspended in SM buffer, 50 mM Tris-Cl, pH 7.5, 100 mM NaCl, 0.2% MgSO₄•7H₂O, and 0.1% gelatin, and purified by a secondary screening following the same procedure. Twenty three strongly hybridizing plaques were identified by the first screening from a total of 3x10⁵ pfu, and fifteen were later purified. DNAs were prepared from two of the purified phage clones (Ausubel et al., (1990) pp. 1.13.4-1.13.8), they were digested with BamHI, ClaI, PstI, and NcoI and prepared for a Southern blot. The blot was hybridized with the SAMS 315 bp PstI fragment probe prepared and used as above. A single positive fragment of clone 1 was identified from the ClaI digestion. Since the ClaI restriction site in the cDNA clone is 843 bp from the 5' end of the full length cDNA, the 2.5 kb ClaI fragment was expected to include about 1.7 kb of DNA upstream of the coding sequence, which was considered sufficient to contain the SAMS promoter.

The 2.5 kb ClaI genomic DNA fragment was cloned into pBluescript™ KS and the DNA insert was sequenced. The 3' end sequence of the genomic DNA fragment was expected to match the 5' end sequence of SAMS cDNA from the 5' end to the ClaI site at position 843. However, comparison of the genomic DNA sequence and the cDNA sequence

revealed that the two sequences have 191 bp of overlapping sequence starting at position 54 and ending at position 245 of the cDNA sequence (SEQ ID NO:1). The sequence of the 2.5 kb genomic DNA clone downstream of the 191 bp overlapping region was determined to be derived from the cloning vector, lambda EMBL3 SP6/T7, which contributed 257 bp of sequence to the 3' end of the 2.5 kb SAMS ClaI fragment including the ClaI cloning site. Therefore, the soybean derived DNA in the 2.5 kb ClaI fragment is described by the 2336 bp DNA sequence shown in SEQ ID NO:2.

The DNA sequence of the genomic DNA in the 191 bp region (from nucleotide 2145 to the end of the sequence) was very similar to, but did not match perfectly, the cDNA sequence; there were six base pair mismatches in this region. This was not surprising, because it was known from the experiments described in Example 3 that there is a small family of SAMS genes in soybean. It was concluded that this genomic clone is not derived from the same gene from which the cDNA used as the probe was transcribed. It was also noted that the 53 bp at the 5' end of the cDNA did not show any similarity to the genomic sequence upstream of the 191 bp overlapping region (Figure 2).

A BLASTN search of the DuPont soybean EST database using the nucleotide sequence from the soybean SAMS genomic DNA upstream of the 191 bp region revealed many cDNA clones that matched a 60 bp region of the genomic DNA from nucleotide 1496 to 1555. The sequence of one such cDNA, designated srr1c.pk002.b21, is shown in SEQ ID NO:3.

The cDNA sequence in SEQ ID NO:3 perfectly matches the genomic sequence in SEQ ID NO:2 from nucleotide 2 to 60 of the cDNA. There follows a region of 591 nucleotides in the genomic DNA that is absent from the cDNA. Then the region from nucleotide 60 to 250 of the cDNA perfectly matches the 191 bp region at the 3' end of the genomic DNA. This indicates the presence of a 591 nucleotide intron in the genomic DNA in the 5' transcribed, but untranslated, region of the SAMS gene. The presence of consensus 5' and 3' splice junctions in the genomic DNA at the exon-intron junctions supports this conclusion. Thus, the 53 bp at the 5' end of the cDNA used as the probe (SEQ ID NO:1) did not match the genomic sequence because the genomic sequence at that position in the alignment was from the intron. However, the 53 bp at the 5' end of the cDNA of SEQ ID NO:1 is very similar to the 60 nucleotides at the 5' end of the cDNA of SEQ ID NO:3, suggesting that the gene from which SEQ ID NO:1 was transcribed also contains an intron at the analogous position.

A 1305 bp SAMS genomic DNA fragment starting at nucleotide 856 and ending at nucleotide 2160 of SEQ ID NO:2: was amplified by PCR from the 2.5 kb ClaI clone. The promoter fragment was amplified from this fragment using primers sam-5 (SEQ ID NO:4) and sam-6 (SEQ ID NO:5) and Pfu DNA polymerase (Stratagene).

CATGCCATGGCTTTATACTTCAAAAAGTGCAC (SEQ ID NO:4)

GCTCTAGATCAAACACATCCAA (SEQ ID NO:5)

5 An XbaI site and an NcoI site were introduced to the 5' end and 3' end, respectively, of the PCR fragment by using these specifically designed primers. The NcoI site includes the ATG start codon of the SAMS coding region. The resulting 1314 bp fragment is shown in SEQ ID NO:6 and includes the SAMS promoter and the translation leader region, which is interrupted by the 591 nucleotide intron.

10 Using PCR amplification procedures and appropriate primers additional SAMS promoter fragments can be produced from the 2336 nucleotide fragment of SEQ ID NO:2. These include, but are not limited to, the three fragments provided in SEQ ID NOs:14, 15 and 16. SEQ ID NO:14 is a 2165 nucleotide sequence of a SAMS promoter DNA fragment which starts at the 5' end of the 2336 nucleotide sequence of SEQ ID NO:2 and ends at the
15 ATG translation start codon of the SAMS protein. SEQ ID NO:15 is a 1574 nucleotide sequence of a SAMS promoter DNA fragment which starts at the 5' end of the 2336 nucleotide sequence of SEQ ID NO:2 and ends at the ATG translation start codon of the SAMS protein, and from which the 591 nucleotide long intron sequence has been removed. SEQ ID NO:16 is a 719 nucleotide sequence of a SAMS promoter DNA fragment which
20 starts at nucleotide 4 of SEQ ID NO:6 and ends at the ATG translation start codon of the SAMS protein, and from which the 591 nucleotide long intron sequence has been removed.

EXAMPLE 5

Expression of the GUS Gene by the SAMS Promoter in *Arabidopsis*

The activity of the soybean SAMS promoter was tested by its ability to express the
25 GUS reporter gene in transgenic *Arabidopsis* plants carrying the SAMS promoter::GUS::3' Nos expression cassette. GUS refers to the *E. coli* β -glucuronidase gene (GUS) (Jefferson, (1987) *Plant Mol. Biol. Rep.* 5:387-405) and 3' Nos refers to the transcription termination region from the nopaline synthase (Nos) gene (Depicker *et al.* (1982) *J. Mol. Appl. Genet.* 1:561-570). The SAMS promoter fragment (SEQ ID NO:6) was digested with XbaI and
30 NcoI and inserted into plasmid pMH40 Δ (SEQ ID NO:17), which contained a 35S promoter::GUS::3' Nos plant expression cassette. The XbaI/NcoI SAMS promoter DNA fragment replaced the 35S promoter of pMH40 Δ , to form the pZSL11 plasmid (Figure 3). The SAMS promoter::GUS::3' Nos DNA fragment (SEQ ID NO:18) was excised from pZSL11 by HindIII and SacI digestion and transferred into the corresponding sites of pBI101
35 (ClonTech) binary vector. The cloned SAMS promoter was sequenced to verify that no sequence error was generated by the PCR amplification.

The SAMS::GUS expression cassette was introduced into wild type *Arabidopsis thaliana* by *Agrobacterium* mediated transformation. *A. thaliana* ecotype columbia were grown in 228 chamber with continuous light and transformed by vacuum infiltration method

using GV3101 *Agrobacteria* (Bent, A. *et al.*, (1994) *Science* 265:1856-1860). Transformed *Arabidopsis* seeds were selected by germination on Murashige and Skoog minimal salt (GIBCO BRL) plus 0.2 % phytigel (Sigma), 1% sucrose, and 100 mg/ml kanamycin. The kanamycin resistant seedlings were transferred into soil and grown in 228 chamber under continuous light.

For histochemical GUS staining, plant tissues were incubated in 0.5% 5-bromo-4-chloro-3-indoxyl- β -D-glucuronic acid (X gluc, Biosynth AG, Switzerland) in 50 mM sodium phosphate, pH 7.0, 10 mM EDTA, 0.5 mM potassium ferricyanide, and 0.5 mM potassium ferrocyanide at 37 $^{\circ}$ C overnight, and then chlorophyll was removed with 75% ethanol. Pictures were taken using a Nikon dissecting microscope. Strong GUS expression was detected in all the parts of the transgenic *Arabidopsis* plants, including flowers (Figure 4A), leaves (Figure 4B), stems (bolt) (Figure 4C), silique coats and developing seeds (Figure 4D-F), developing embryos (Figure 4G), and seedlings (not shown). The GUS staining on leaves and silique coats was uniform with all the veins and mesophyll tissues similarly stained, while staining on flowers and stems was not uniform. Although some seeds were not stained for GUS activity due to genetic segregation, the funiculi that connected these seeds to the silique coat stained positively for GUS activity (Figure 4G). These results indicated that the soybean SAMS promoter was a constitutive promoter and was able to function in heterologous plant.

The GUS activities of the transgenic *Arabidopsis* plants were further analyzed by a fluorometric assay. For fluorescence analysis, plant tissues were ground in microfuge tubes with extraction buffer, 50 mM phosphate buffer, pH 7.0, 10 mM EDTA, 0.1% Triton X-100, 0.1% N-lauroyl sarcosine, and 10 mM β -mercaptoethanol, to homogeneity. The samples were centrifuged at 14,000 rpm for 10 minutes, and aliquots of the supernatant were used to determine protein concentrations by the Bradford method (Bio-Rad, Hercules, CA) using 96 well microtiter plates read with a kinetic microplate reader (Molecular Devices, Sunnyvale, CA). The β -glucuronidase activities were analyzed by standard protocol (Jefferson *et al.*, (1987) *EMBO J.* 6:3901-3907) using 96 well microtiter plates read with Cytofluor multiwell plate reader (PerSeptive Biosystems, Framingham, MA). Data were entered into a Microsoft Excel spread sheet and analyzed. Triple samples of flower, leaf, stem, silique coat, young seed (white), medium seed (light green), old seed (dark green), and dry seed from six plants were analyzed. The soybean SAMS promoter was active in all the tissues analyzed (Figure 5). Promoter activity varied among the six lines, as is typically seen among plant transformants. The basic expression patterns were similar among all the lines, and the average SAMS promoter activity was comparable to that of the 35S promoter (Battraw and Hall, (1990) *Plant Mol. Biol.* 15:527-538; Jefferson *et al.*, (1987) *EMBO J.* 6:3901-3907; Atanassova *et al.*, (1998) *Plant Mol. Biol.* 37:275-285; Holtorf *et al.*, (1995) *Plant Mol. Biol.* 29:637-646; Wilpink *et al.*, (1995) *Plant Mol. Biol.* 28:949-955). The

SAMS promoter was very active in developing seeds, especially in early and medium stages of development, and the GUS specific activities are in the range of 5-40 pmole 4-Mu (4-methylumbelliferone) per microgram protein per minute, which are comparable to many strong promoters (Atanassova *et al.*, (1998) *Plant Mol. Biol.* 37:275-285; Comai *et al.*, (1990) *Plant Mol. Biol.* 15:373-381; Holtorf *et al.*, (1995) *Plant Mol. Biol.* 29:637-646; Wilmink *et al.*, (1995) *Plant Mol. Biol.* 28:949-955).

EXAMPLE 6

Expression of GUS Gene by SAMS Promoter in Corn

In order to test whether the dicot SAMS promoter also worked in monocot plants, pZSL11 was introduced into corn leaf discs and callus by gene bombardment for transient gene expression assay using the biolistic particle delivery system PDS-1000/He (Bio Rad, Hercules, CA). The pMH40Δ plasmid DNA (as set forth in SEQ ID NO:17), which contained the 35S promoter and GUS reporter gene, was also introduced into corn callus and leaf discs by gene bombardment to serve as a positive control vector. After incubation overnight at 37°, bombarded tissues were stained for GUS activity. GUS expression was demonstrated by the blue spots on both the callus (Figure 6A) and leaf discs (Figure 6B) bombarded with pZSL11. As expected, the positive control 35S::GUS cassette was also expressed in both callus and leaf discs (Figure 6C, D).

EXAMPLE 7

Expression of Methionine Biosynthesis Genes by SAMS Promoter

The SAMS promoter was fused to two soybean cDNAs, one encoding ATP sulfurylase (ATPS) and a second encoding cystathionine-γ-synthase (CGS1). The soybean ATPS and CGS1 cDNAs were isolated from soybean embryo cDNA libraries using the same procedures as described in Example 1 and Example 2 for isolation of soybean SAMS cDNAs. The coding regions and the 3' untranslated region (UTR) of soybean ATPS and CGS1 genes were inserted into pZSL11 replacing the GUS gene. The resulting SAMS promoter::ATPS and SAMS promoter::CGS1 expression cassettes, SEQ ID NO:19 and SEQ ID NO:20, respectively, were inserted into binary vectors for *Arabidopsis* transformation and transformation was performed as described in Example 5. Transgenic *Arabidopsis* plants with soybean ATPS and CGS1 genes controlled by the SAMS promoter were analyzed by PCR for the presence of the transgenes and by RT-PCR for expression of the transgenes. Genomic DNA used for PCR analysis was prepared from *Arabidopsis* siliques and leaves using 7 M urea, 1.5 M NaCl, 50 mM Tris, pH 8.0, 20 mM EDTA, and 1% N-lauroyl-sarcosine, followed by phenol extraction and ethanol precipitation. Primer sam-9 (SEQ ID NO:7) which is specific to SAMS promoter, and primers specific to the target genes, atps-1 (SEQ ID NO:8) for the ATPS gene and cgs-8 (SEQ ID NO:9) for the CGS1 gene were used in PCR with Taq DNA polymerase (GIBCO BRL) to detect the existence of SAMS::ATPS and SAMS::CGS1 in transgenic *Arabidopsis* plants.

TTCGAGTATAGGTCACAATAGG (SEQ ID NO:7)
 CTTCGCTGAGGACATGGAC (SEQ ID NO:8)
 5 GAGTTGTCGCTGTTGTTTCGAC (SEQ ID NO:9)

RNA samples used for RT-PCR were prepared with Trizol™ Reagent (GIBCO BRL). Antisense primers atps-4 (SEQ ID NO:10) and cgs-10 (SEQ ID NO:11) were used in reverse transcription reactions with SuperscriptII™ RT (GIBCO BRL) following the vendor's instruction.

AACACAGCATCCGCATTGCG (SEQ ID NO:10)
 15 AGGAGTGCAGAATCAGATCAG (SEQ ID NO:11)

The first strand cDNAs were used in PCR with primer pairs atps-3 (SEQ ID NO:12) and atps-4 (SEQ ID NO:10) for SAMS::ATPS transgenic plants, and cgs-9 (SEQ ID NO:13) and cgs-10 for SAMS::CGS1 transgenic plants. PCR and RT-PCR products were resolved by agarose gel electrophoresis.

GCTGATCGAACCAGATGGAG (SEQ ID NO:12)
 25 CTGTACAGTTAAACAGTAGTTCT (SEQ ID NO:13)

All ten SAMS::CGS1 transgenic *Arabidopsis* harbored the SAM::CGS1 expression cassette as revealed by PCR with SAMS::CGS1-specific primers (Figure 7A). It was also revealed by the same analysis that all the ten SAMS::ATPS transgenic *Arabidopsis* plants contained the SAMS::ATPS expression cassette (Figure 7A). RT-PCR analysis detected CGS1 transcripts and ATPS transcripts, respectively, in most of the transgenic plants (Figure 7B). This shows that the SAMS promoter is capable of driving expression of a variety of different genes in most or all cell types in transformed plants.

EXAMPLE 8

Induction of SAMS Promoter Activity by Methionine

Since SAMS catalyzes the reaction to synthesize S-adenosyl-L-methionine from methionine and ATP, free methionine levels might regulate SAMS promoter activity. To see if SAMS promoter is regulated by external methionine, the SAMS::GUS transgenic *Arabidopsis* seeds were germinated in the presence of either H₂O, 1x Murashige and Skoog salt (GIBCO BRL), 0.01 mM methionine (Sigma), or 1 mM methionine. Ten days old seedlings from ten independent transgenic lines were analyzed for GUS activity according to the protocol described in Example 5. GUS activity for each treatment, in the order given above, for each transgenic line is shown in Figure 8. All lines responded similarly to the different treatments. Compared to the control of H₂O treatment, SAMS activity was

induced more than two-fold by 0.01 mM free methionine and inhibited about 40% on average by 1x MS salt. The induction effect of SAMS promoter by 1 mM methionine was less than that by 0.01 mM methionine, probably due to a toxic effect of the high methionine concentration; this toxic effect was indicated by the smaller sizes and shorter roots of the seedlings grown in the presence of 1 mM methionine. The toxic effect of high levels of methionine was even more apparent at 10 mM free methionine, since only a few *Arabidopsis* seeds were able to germinate and none survived in the presence of 10 mM free methionine.

CLAIMS

What is claimed is:

1. An isolated nucleic acid fragment comprising a promoter wherein said promoter consists essentially of the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15, or 16 or said promoter consists essentially of a fragment or subfragment that is substantially similar and functionally equivalent to the nucleotide sequence set forth in SEQ ID NOs:6, 14, 15, or 16.
2. A chimeric gene comprising at least one heterologous nucleic acid fragment operably linked to the promoter of Claim 1 or Claim 10.
3. A plant containing the chimeric gene of Claim 2.
4. The plant of Claim 3 wherein said plant is a monocot selected from the group consisting of corn, rice, wheat, barley and palm.
5. The plant of Claim 3 wherein said plant is a dicot selected from the group consisting of *Arabidopsis*, soybean, oilseed *Brassica*, peanut, sunflower, safflower, cotton, tobacco, tomato, potato, and cocoa.
6. The plant of claim 5 wherein said plant is soybean.
7. Seeds of the plants of Claims 3, 4, 5, or 6.
8. A method of increasing or decreasing the expression of at least one heterologous nucleic acid fragment in a plant cell which comprises:
 - (a) transforming a plant cell with the chimeric gene of Claim 2;
 - (b) growing fertile mature plants from the transformed plant cell of step (a);
 - (c) selecting plants containing a transformed plant cell wherein the expression of the heterologous nucleic acid fragment is increased or decreased.
9. The method of Claim 8 wherein the plant is a monocot selected from the group consisting of corn, rice, wheat, barley and palm.
10. The method of Claim 9 wherein the plant is a dicot selected from the group consisting of *Arabidopsis*, soybean, oilseed *Brassica*, peanut, sunflower, safflower, cotton, tobacco, tomato, potato, and cocoa.
11. The method of Claim 10 wherein the plant is soybean.
12. An isolated nucleic acid fragment comprising a constitutive plant SAMS promoter.

TITLE

S-ADENOSYL-L-METHIONINE SYNTHETASE PROMOTER
AND ITS USE IN EXPRESSION OF TRANSGENIC GENES IN PLANTS

ABSTRACT OF THE DISCLOSURE

5 A constitutive plant S-adenosyl-L-methionine synthetase (SAMS) promoter and
subfragments thereof and their use in promoting the expression of one or more heterologous
nucleic acid fragments in plants are described.

10

15

20


25

30

35


LMC/bjm

FIGURE 1 A



Soybean genomic DNA Southern blot hybridized
with the SAM5 cDNA probe.

FIGURE 1B



Soybean genomic DNA Southern blot hybridized
with the SAM5 promoter probe.

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100
1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	

AGCCAAGCCCCACTCAACCACCACACCCTCTCTGCTCTTCTTCTACCTTTCAAGTTTTTAAAGTATTA

GCTGTCCTCGACGCTTGCCCTGAACAGGACCCAGACAGCAAGGTTGCCTGCGAAACATGCACCAAGACCAACTTGGTCAT
GCTGTCCTCGACGCTTGCCCTGAACAGGACCCAGACAGCAAGGTTGCCTGCGAAACATGCACCAAGACCAACTTGGTCAT

GGTCTTCGGAGAGATC
: : : : : : : : : : : : : : : :
GGTCTTCGGAGAGATC

FIGURE 3




FIGURE 4





FIGURE 5



22






FIGURE 7





Induction of SAMS promoter activity by free methionine. Seeds of ten transgenic *Arabidopsis* lines transformed with SAMS::GUS construct were germinated on filter papers soaked with H₂O, 1x Murashige and Skoog salt, 0.01 mM, and 1 mM methionine. Ten days old seedlings were harvested and assayed for GUS activity. The solid bar and hollow bar indicate, respectively, the average and the standard variation of three samples for treatment.

L R S Y M O P



DECLARATION and POWER OF ATTORNEY

As a below-named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**S-ADENOSYL-L-METHIONINE SYNTHETASE PROMOTER AND ITS USE IN EXPRESSION OF
TRANSGENIC GENES IN PLANTS**

the specification of which is attached hereto unless the following box is checked:

☐ was filed on _____ as U.S. Application No. _____ or PCT International Application No. _____ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Application No.	Country	Filing Date	Priority Claimed (Yes/No)
-----------------	---------	-------------	---------------------------

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States Provisional Application(s) listed below.

U.S. Provisional Application No.

60/113,045

U.S. Filing Date

12/21/98

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International Application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application or PCT International Application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is known to me to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application No.	Filing Date	Status (patented, pending or abandoned)
-----------------	-------------	---

POWER OF ATTORNEY: I hereby appoint the following attorney(s) and/or agent(s) the power to prosecute this application and transact all business in the Patent and Trademark Office connected therewith:

Name:	LYNNE M. CHRISTENBURY	Registration No.:	30,971
Send correspondence and direct telephone calls to:	LYNNE M. CHRISTENBURY	E. I. du Pont de Nemours and Company Legal - Patents Wilmington, DE 19898, U.S.A.	Tel. No. (302) 992-5481 Fax No. (302) 892-7949

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

INVENTOR(S)

Full Name of Inventor	Last Name FALCO	First Name SAVERIO	Middle Name CARL
	Signature (please sign full name):		Date:
Residence & Citizenship	City ARDEN	State or Foreign Country DELAWARE	Country of Citizenship U.S.A.
Post Office Address	Post Office Address 1902 MILLERS ROAD	City ARDEN	State or Country DELAWARE
			Zip Code 19810
Full Name of Inventor	Last Name LI	First Name ZHONGSEN	Middle Name
	Signature (please sign full name):		Date:
Residence & Citizenship	City HOCKESSIN	State or Foreign Country DELAWARE	Country of Citizenship PEOPLE'S REPUBLIC OF CHINA
Post Office Address	Post Office Address 24 FOREST CREEK DRIVE	City HOCKESSIN	State or Country DELAWARE
			Zip Code 19707

☐ Additional Inventors are being named on separately numbered sheets attached hereto.

SEQUENCE LISTING

<110> FALCO, SAVERIO CARL
LI, ZHONGSEN

<120> S-ADENOSYL-L-METHIONINE SYNTHETASE PROMOTER AND
ITS USE IN EXPRESSION OF TRANSGENIC GENES IN PLANTS

<130> BB1205 US NA

<140>

<141>

<150> 60/113,045

<151> 1998-12-21

<160> 16

<170> Microsoft Office 97

<210> 1

<211> 1518

<212> DNA

<213> Glycine max

<400> 1

```

agccaagccc cactcaacca ccacaccact ctctctgctc ttcttctacc tttcaagttt 60
ttaaagtatt aagatggcag agacattcct atttacctca gagtcagtga acgagggaca 120
ccctgacaag ctctgcgacc aaatctccga tgctgtcctc gacgcttgcc ttgaacagga 180
cccagacagc aaggttgccct gcgaaacatg caccaagacc aacttggtca tggctctcgg 240
agagatcacc accaaggcca acgttgacta cgagaagatc gtgctgaca cctgcaggaa 300
catcggcttc gtctcaaacg atgtgggact tgatgctgac aactgcaagg tccttgtaaa 360
cattgagcag cagagccctg atattgccca ggggtgtgcac ggccacctta ccaaagacc 420
cgaggaaatc ggtgctggag accagggtca catgtttggc tatgccacgg acgaaacccc 480
agaattgatg ccattgagtc atgttcttgc aactaaactc ggtgctcgtc tcaccgaggt 540
tcgcaagaac ggaacctgcc catggttgag gcctgatggg aaaacccaag tgactgttga 600
gtattacaat gacaacggtg ccatggttcc agttcgtgtc cacactgtgc ttatctccac 660
ccaacatgat gagactgtga ccaacgcagc aattgcagct gacctcaagg agcatgtgat 720
caagccgggtg atcccggaga agtaccttga tgagaagacc attttccact tgaacccctc 780
tgcccgtttt gtcattggag gtccctcacgg tgatgctggg ctacccggcc gcaagatcat 840
catcgatact tacggaggat ggggtgctca tgggtgggtg gctttctccg ggaaggatcc 900
caccaagggt gataggagt gtgcttacat tgtgagacag gctgctaaga gcattgtggc 960
aagtggacta gccagaaggt gcattgtgca agtgtcttat gccattggtg tgcccagacc 1020
tttgtctgtc tttgttgaca cctatggcac cgggaagatc catgataagg agattctcaa 1080
cattgtgaag gagaactttg atttcaggcc cggtatgata tccatcaacc ttgatctcaa 1140
gaggggtggg aataacaggt tcttgaagac tgctgcataa ggacacttcg gcagagagga 1200
ccctgacttc acatgggaag tggtaagacc cctcaagtgg gagaaggcct aaggccattc 1260
attccactgc aatgtgctgg gagtttttta gcgttgccct tataatgtct attatccata 1320
actttccacg tcccttgctc tgtgtttttc tctcgtcgtc ctccctctat tttgtttctc 1380
ctgcctttca tttgtaattt tttacatgat caactaaaaa atgtactctc tgttttccga 1440
ccattgtgtc tcttaatatc agtatcaaaa agaattgttc aagttaaaaa aaaaaaaaaa 1500
aaaaaaaaaa aaaaaaaaaa
1518

```

<210> 2

<211> 2336

<212> DNA

<213> Glycine max

```

<400> 2
atcgatagag acatgttatt cacaaaccat aaaatgatgg ctaaaattgg tgtgattgga 60
acgatatctg tttattatga tttcagggcg caaaaatgcg agtacttaat aaaattttac 120
atttaaatta gaattttttt tatcaataaa tattaattta ttagttttat tagaaatatt 180
aattagaaaa ttttgaatcc ccgattttct ctccttttct tcgctattca tcattttcta 240
accaaaacca tcttatatgt tcttcaaatt agaacttgaa attattaatt ataattaaac 300
tgaaaacaat ttggtatcaa ttcataataca tgcttagtaa taaaatgcga taattaattg 360
ataaatctgc aaaagatttt acaaatatct ttcagaaaaa attaataaca aattttgtcg 420
ttttcatggg gttgggtctga ggaggatttg gcactataga actctcctac ggaccattct 480
ttgcacttca actaaacgat ggtcagaatt ggtggggatt ttatattcaa gcataatccct 540
ttcaaaactt cctacttact tcgtgcgttc ggtaatcggg aacattagac tttcaaaatc 600
atttttaacc cctaaacagt aaatttgaag gacaaaaata atatttttca aatttgatag 660
actatttttt ttttgtaatt tgacgaacca aaaccagatt tatcctgaat tttaggaacc 720
acagatgtaa ctaaaccaat atttatttat tttctaaaaa aaaatttcat gccagcatgc 780
ctcagcccat gaaaaaaaacc ttataaaaaat atctacacat tgaccattga aaagttcggt 840
ctcccatggg taaccagatc aaactcacat ccaaacataa catggatac tccttaccac 900
tcataactaat tattttgggt taaatattaa tcattatttt taagatatta attaagaaat 960
taaaagattt tttaaaaaaa tgtataaaaat tatattatct atgatttttc atacatttga 1020
ttttgataat aaatatattt tttttaattt cttaaaaaat gttgcaagac acttattaga 1080
catagtcttg ttctgtttac aaaagcattc atcatttaat acattaaaaa atattttaata 1140
ctaacagtag aatcttcttg tgagtgggtg gggagtaggc aacctggcat tgaaacgaga 1200
gaaagagagt cagaaccaga agacaaataa aaagtatgca acaaacaaat caaaatcaaa 1260
gggcaaaggc tgggggttggc tcaattgggt gctacattca attttcaact cagtcaacgg 1320
ttgagattca ctctgacttc cccaatctaa gccgcggatg caaacggttg aatctaacc 1380
acaatccaat ctogttactt aggggctttt ccgtcattaa ctacccctg ccaccgggtt 1440
tcctataaaa ttggaactca atgctccct ctaaactcgt atcgcttcag agttgagacc 1500
aagacacact cgttcatata tctctctgct cttctcttct cttctacctc tcaagggtact 1560
tttcttctcc ctctaccaa tcttagattc cgtggttcaa tttcggatct tgcacttctg 1620
gtttgctttg ccttgctttt tctcaactg ggtccatcta ggatccatgt gaaactctac 1680
tctttcttta atatctgcgg aatacgcgtt ggactttcag atctagtcga aatcatttca 1740
taattgcctt tctttctttt agcttatgag aaataaaaatc attttttttt atttcaaaat 1800
aaaccttggg ccttggtgctg actgagatgg ggtttggtga ttacagaatt ttagcgaatt 1860
ttgtaattgt acttgtttgt ctgtagtttt gttttgtttt cttgtttctc atacattcct 1920
taggcttcaa ttttattcga gtataggcca caataggaat tcaaactttg agcaggggaa 1980
ttaatccctt ccttcaaact cagtttggtt gtatatatgt ttaaaaaatg aaacttttgc 2040
tttaaatctt attataactt tttttatggc aaaaattttt gcatgtgtct ttgctctcct 2100
gttgtaaat tactgtttag gtactaactc taggcttggt gtgcagtttt tgaagtataa 2160
agatggcaga gacattccta ttcacctcgg agtcagtgaa cgaggacac cctgataagc 2220
tctgcgacca aatctccgat gctgtcctcg acgcttgctt cgaacaggac ccagacagca 2280
aggttgcttg cgaaacatgc accaagacca acttgggtcat ggtcttcgga gagatc 2336

```

```

<210> 3
<211> 522
<212> DNA
<213> Glycine max

```

```

<220>
<221> unsure
<222> (405)

```

```

<220>
<221> unsure
<222> (509)

```

```

<220>
<221> unsure
<222> (515)

```

```

<400> 3
gaccaagaca cactogttca tataatctctc tgctcttctc ttctcttcta cctctcaagt 60
ttttgaagta taaagatggc agagacattc ctattcacct cggagtcagt gaacgagggga 120
caccctgata agctctgcga ccaaattctcc gatgctgtcc tcgacgcttg cctcgaacag 180
gacccagaca gcaagggttg ctgcgaaaca tgcaccaaga ccaacttggt catggtcttc 240
ggagagatca ccaaccaagg caacgttgac tacgagaaga tcgtgcgtga cacctgcagg 300
agcatcggtt tcatctcaaa cgatgtggga cttgatgctg acaactgcaa ggtccttgta 360
aacattgagc agcagagccc tgatattgcc cagggcgtgc acggncacct taccaaaaga 420
cctgaagaaa ttggcgctgg tgaccaaggt cacatgtttg gctatgccac tgatgaaacc 480
ccaaaattca tgccattgag tcatgttctt gcaancaagc tc 522

```

```

<210> 4
<211> 32
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: PCR Primer

```

```

<400> 4
catgccatgg cttttatactt caaaaactgc ac 32

```

```

<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence

```

```

<220>
<223> Description of Artificial Sequence: PCR Primer

```

```

<400> 5
gctctagatc aaactcacat ccaa 24

```

```

<210> 6
<211> 1314
<212> DNA
<213> Glycine max

```

```

<400> 6
tctagatcaa actcacatcc aaacataaca tggatatctc cttaccaatc atactaatta 60
ttttgggtta aatattaatc attattttta agatattaat taagaaatta aaagattttt 120
taaaaaaatg tataaaatta tattattcat gatttttcat acatttgatt ttgataataa 180
atatattttt tttaatttct taaaaaatgt tgcaagacac ttattagaca tagtcttggt 240
ctgttttaca aagcattcat catttaatac attaaaaaat atttaatact aacagtagaa 300
tcttcttggt agtggtgtgg gagtaggcaa cctggcattg aaacgagaga aagagagtca 360
gaaccagaag acaaataaaa agtatgcaac aaacaaatca aaatcaaagg gcaaaggctg 420
gggttggttc aattgggttg tacattcaat ttccaactca gtcaacgggt gagattcact 480
ctgacttccc caatctaagc cgcggatgca aacgggtgaa tctaaccac aatccaatct 540
cgttacttag gggcttttcc gtcattaact caaccctgcc acccggttc cctataaatt 600
ggaactcaat gtcctcctct aaactcgtat cgcttcagag ttgagaccaa gacacactcg 660
ttcatatata tctctgctct tctcttctct tctacctctc aaggtacttt tcttctccct 720
ctaccaaate ctagattccg ttggtcaatt tcggatcttg cacttctggt ttgctttgcc 780
ttgctttttc ctcaactggg tccatctagg atccatgtga aactctactc tttctttaat 840
atctgcgga tgcggttg actttcagat ctagtgcgaa tcatttcata attgcctttc 900
tttcttttag cttatgagaa ataaaatcat tttttttat ttcaaaataa accttgggccc 960
ttgtgctgac tgagatgggg tttggtgatt acagaatttt agcgaatttt gtaattgtac 1020
ttgtttgtct gtagttttgt tttgttttct tgtttctcat acattcctta ggcttcaatt 1080
ttattcgagt ataggtcaca ataggaattc aaactttgag caggggaatt aatcccttcc 1140
ttcaaatcca gtttgtttgt atatatgttt aaaaaatgaa acttttgctt taaattctat 1200

```

tataactttt tttatggcaa aaatttttgc atgtgtcttt gctctcctgt tgtaaattta 1260
ctgttttaggt actaactcta ggcttggtgt gcagtttttg aagtataacc atgg 1314

<210> 7
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 7
ttcgagtata ggtcacaata gg 22

<210> 8
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 8
cttcgctgag gacatggac 19

<210> 9
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 9
gagttgtcgc tgttgttcga c 21

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 10
aacacagcat ccgcattgcg 20

<210> 11
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 11
aggagtgcag aatcagatca g 21

<210> 12
<211> 20

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 12
gctgatcgaa ccagatggag 20

<210> 13
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: PCR Primer

<400> 13
ctgtacagtt aaacagtagt tct 23

<210> 14
<211> 2165
<212> DNA
<213> Glycine max

<400> 14
atcgatagag acatgttatt cacaaacat aaaatgatgg ctaaaattgg tgtgattgga 60
acgatatctg tttattatga tttcagggcg caaaaatgcg agtacttaat aaaatttttac 120
atttaaatta gaatTTTTTT tatcaataaa tattaattta ttagttttat tagaaatatt 180
aattagaaaa ttttgaatcc cggatttctc ctccttttct tcgctattca tcatttttcta 240
accaaaccac tcttatatgt tcttcaaatt agaacttgaa attattaatt ataattaaac 300
tgaaaacaat ttggtatcaa ttcatatata tgcttagtaa taaaatgcga taattaattg 360
ataaatctgc aaaagatttt acaaatatct ttcagaaaaa attaataaca aattttgtcg 420
ttttcatggt gttggtctga ggaggtttg gcactataga actctcctac ggaccattct 480
ttgcacttca actaaacgat ggtcagaatt ggtggggatt ttatattcaa gcatatccct 540
ttcaaaactt cctacttact tcgtgcgttc ggtaatcggg aacattagac tttcaaaatc 600
atttttaacc cctaaacagt aaatttgaag gacaaaaata atatttttca aatttgatag 660
actatttttt ttttghtaatt tgacgaacca aaaccagatt tatcctgaat tttaggaacc 720
acagatgtaa ctaaaccaat atttatttat tttctaaaaa aaaatttcat ggcagcatgc 780
ctcagcccat gaaaaaaacc ttataaaaaa atctacacat tgaccattga aaagtctcgtt 840
ctcccatggg taaccagatc aaactcacat ccaaacataa catggatata tccttaccac 900
tcatactaata ttttttgggt taaatattaa tcattatttt taagatatta attaagaaat 960
ttaaagattt tttaaaaaaa ttataaaaaa tatattattc atgatttttc atacatttga 1020
ttttgataat aaatatattt tttttaattt cttaaaaaat gttgcaagac acttattaga 1080
catagtcttg ttctgtttac aaaagcattc atcatttaat acattaaaaa atatttaata 1140
ctaacagtag aatcttcttg tgagtgggtg gggagtaggc aacctggcat tgaaacgaga 1200
gaaagagagt cagaaccaga agacaaataa aaagtatgca acaaacaaat caaaatcaaa 1260
gggcaaaggc tggggttggc tcaattgggt gctacattca attttcaact cagtcaacgg 1320
ttgagattca ctctgacttc cccaatctaa gccgcggatg caaacggttg aatctaacc 1380
acaatccaat ctcgttactt aggggctttt ccgtcattaa ctcacccctg ccaccggtt 1440
tcctataaaa ttggaactca atgctccctt ctaaactcgt atcgcttcag agttgagacc 1500
aagacacact cgttcatata tctctctgct cttctcttct cttctacctc tcaagggtact 1560
tttcttctcc ctctaccaa tcttagattc cgtggttcaa tttcggatct tgcacttctg 1620
gtttgctttg ccttgctttt tcttcaactg ggtccatcta ggatccatgt gaaactctac 1680
tctttcttta atatctgcgg aatacgcgtt ggactttcag atctagtcga aatcatttca 1740
taattgcctt tctttctttt agcttatgag aaataaaatc attttttttt atttcaaaat 1800
aaaccttggg ccttggtgctg actgagatgg ggttgggtga ttacagaatt tttagcgaatt 1860
ttgtaattgt acttgtttgt ctgtagtttt gttttgtttt cttgttttct atacattcct 1920
taggcttcaa ttttattcga gtataggtea caataggaat tcaaaccttg agcaggggaa 1980
ttaatccctt ctttcaaatc cagtttgttt gtatatatgt ttaaaaaatg aaacttttgc 2040

ttttaaattct	attataaactt	tttttatggc	aaaaattttt	gcatgtgtct	ttgctctcct	2100
gttgtaaatt	tactgttttag	gtactaactc	taggcttggt	gtgcagtttt	tgaagtataa	2160
agatg						2165

<210> 15
 <211> 1574
 <212> DNA
 <213> Glycine max

<400> 15						
atcgatagag	acatgttatt	cacaaacccat	aaaatgatgg	ctaaaattgg	tgtgattgga	60
acgatatctg	tttattatga	tttcagggcg	caaaaatgcg	agtacttaat	aaaattttac	120
atttaaatta	gaattttttt	tatcaataaa	tattaattta	ttagttttat	tagaaatatt	180
aattagaaaa	ttttgaatcc	ccgattttct	ctccttttct	tcgctattca	tcattttcta	240
accaaaccaa	tcttatatgt	tcttcaaatt	agaacttgaa	attattaatt	ataattaaac	300
tgaaaacaat	ttgggtatcaa	ttcatatata	tgcttagtaa	taaaatgcga	taattaattg	360
ataaatctgc	aaaagatttt	acaaatatct	ttcagaaaaa	attaataaca	aattttgtcg	420
ttttcatggg	gttgggtctga	ggaggatttg	gcactataga	actctcctac	ggaccattct	480
ttgcacttca	actaaacgat	ggtcagaatt	ggtaggggatt	ttatattcaa	gcatatccct	540
ttcaaaaactt	cctaacttact	tcgtgcgttc	ggtaatcggg	aacatttagac	tttcaaaatc	600
attttttaacc	cctaaacagt	aaatttgaag	gacaaaaata	atatttttca	aatttgatag	660
actatttttt	ttttgttaatt	tgacgaacca	aaaccagatt	tatcctgaat	tttaggaacc	720
acagatgtaa	ctaaaccaat	atttatttat	tttctaaaaac	aaaatttcat	ggcagcatgc	780
ctcagcccat	gaaaaaaaacc	ttataaaaaat	atctacacat	tgaccattga	aaagtctggt	840
ctcccatggg	taaccagatc	aaactcacat	ccaaacataa	catggatatc	tccttaccac	900
tcatactaata	tattttgggt	taaatattaa	tcattatttt	taagatatta	attaagaaat	960
taaaagatttt	tttataaaaaa	tgtataaaaat	tatattattc	atgatttttc	atacatttga	1020
ttttgataat	aaatatattt	tttttaattt	cttaaaaaat	gttgcaagac	acttattaga	1080
catagtcttg	ttctgtttac	aaaagcattc	atcattttaat	acattaaaaa	atatttaata	1140
ctaacagtag	aatcttcttg	tgagtgggtg	gggagtaggc	aacctggcat	tgaaacgaga	1200
gaaagagagt	cagaaccaga	agacaaaata	aaagtatgca	acaaacaaat	caaaatcaaa	1260
gggcaaaggc	tgggggtggc	tcaattgggt	gtacatttca	attttcaact	cagtcaacgg	1320
ttgagattca	ctctgacttc	cccaatctaa	gcgcgggatg	caaacgggtg	aatctaacc	1380
acaatccaat	ctcgttactt	aggggctttt	cogtcattaa	ctcaccctcg	ccaccctgtt	1440
tccctataaaa	ttggaactca	atgctccctc	ctaaactcgt	atcgtctcag	agttgagacc	1500
aagacacact	cgttcatata	tctctctgct	cttctcttct	cttctacctc	tcaagttttt	1560
gaagtataaaa	gatg					1574

<210> 16
 <211> 719
 <212> DNA
 <213> Glycine max

<400> 16						
agatcaaaact	cacatccaaa	cataacatgg	atatctcctt	accaatcata	ctaattattt	60
tgggttaaat	attaatcatt	attttttaaga	tattaattaa	gaaattaaaa	gattttttta	120
aaaaatgtat	aaaattatat	tattcatgat	ttttcatata	tttgattttg	ataataaata	180
tatttttttt	aatttcttaa	aaaatgttgc	aagacactta	ttagacatag	tcttgttctg	240
tttacaaaaag	cattcatcat	ttaatacatt	aaaaaatatt	taataactaac	agtagaatct	300
tcttgtgagt	ggtgtgggag	taggcaacct	ggcattgaaa	cgagagaaaag	agagtcagaa	360
ccagaagaca	aataaaaaag	atgcaacaaa	caaatcaaaa	tcaaagggca	aaggctggg	420
ttgggtcaat	tgggttgctac	attcaatttt	caactcagtc	aacgggttgag	attcactctg	480
acttccccaa	tctaagccgc	ggatgcaaac	ggttgaatct	aaccacacat	ccaatctcgt	540
tacttagggg	cttttccgtc	attaactcac	ccctgccacc	cggtttccct	ataaattgga	600
actcaatgct	cccctctaaa	ctcgtatcgc	ttcagagttg	agaccaagac	acactcgttc	660
atatatctct	ctgctcttct	cttctcttct	acctctcaag	tttttgaagt	ataaagatg	719

<210> 17
 <211> 6975

<212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:plasmid

<400> 17
 gaatatgcat cactagtaag ctttgctcta gaggatccaa ttccaatccc acaaaaatct 60
 gagcttaaca gcacagttgc tcctctcaga gcagaatcgg gtattcaaca ccctcatatc 120
 aactactacg ttgtgtataa cgggtccacat gccgggtatat acgatgactg gggttgtaca 180
 aaggcggcaa caaacggcgt tcccggagtt gcacacaaga aatttgccac tattacagag 240
 gcaagagcag cagctgacgc gtacacaaca agtcagcaaa cagacaggtt gaacttcac 300
 cccaaaggag aagctcaact caagcccaag agctttgcta aggccctaac aagccacca 360
 aagcaaaaag cccactggct cacgctagga accaaaaggc ccagcagtga tccagcccca 420
 aaagagatct cctttgcccc ggagattaca atggacgatt tcctctatct ttacgatcta 480
 ggaaggaagt tcgaagggtga aggtgacgac actatgttca ccactgataa tgagaagggt 540
 agcctcttca atttcagaaa gaatgctgac ccacagatgg ttagagaggc ctacgcagca 600
 ggtctcatca agacgatcta cccgagtaac aatctccagg agatcaaata ccttcccaag 660
 aagggttaaag atgcagtcaa aagattcagg actaattgca tcaagaacac agagaaagac 720
 atattttctca agatcagaag tactattcca gtatggacga ttcaaggctt gcttcataaa 780
 ccaaggcaag taatagagat tggagtctct aaaaaggtag ttccactactga atctaaggcc 840
 atgcatggag tctaagattc aaatcgagga tctaacagaa ctcgccgtga agactggcga 900
 acagttcata cagagtcttt tacgactcaa tgacaagaag aaaatcttcg tcaacatggt 960
 ggagcacgac actctgggtct actccaaaaa tgtcaaagat acagtctcag aagaccaaag 1020
 ggctattgag acttttcaac aaaggataat ttcgggaaac ctcctcggat tccattgccc 1080
 agctatctgt cacttcatcg aaaggacagt agaaaaggaa ggtggctcct acaaatgcca 1140
 tcattgcgat aaaggaaaag ctatcattca agatgcctct gccgacagtg gtcccaaaga 1200
 tggaccccca cccacgagga gcatcgtgga aaaagaagac gttccaacca cgtcttcaaa 1260
 gcaagtggat tgatgtgaca tctccactga cgtaagggat gacgcacaat cccactatcc 1320
 ttcgcaagac ccttctctta tataaggaag ttcatttcat ttggagagga cacgctcgag 1380
 ctcatttctc tattacttca gccataacaa aagaactctt ttctcttctt attaaacat 1440
 ggtacgtcct gtagaaaacc caaccctgta aatcaaaaaa ctcgacggcc tgtgggcatt 1500
 cagtctggat cgcgaaaact gtggaattga tcagcgttgg tgggaaagcg cgttacaaga 1560
 aagccgggca attgctgtgc caggcagttt taacgatcag ttcgccgatg cagatattcg 1620
 taattatgcy ggcaacgtct ggtatcagcg ogaagtcttt ataccgaaag gttgggcagg 1680
 ccagcgtatc gtgctgcgtt tcgatgcggt cactcattac ggcaaagtgt ggggtcaataa 1740
 tcaggaagtg atggagcatc agggcggcta tacgccattt gaagccgatg tcacgccgta 1800
 tgttattgcc gggaaaagtg tacgtatcac cgtttgtgtg aacaacgaac tgaactggca 1860
 gactatcccg ccgggaatgg tgattaccga cgaaaacggc aagaaaaagc agtcttactt 1920
 ccatgatttc tttaactatg ccggaatcca tcgcagcgta atgctctaca ccacgccgaa 1980
 cacctgggtg gacgatatca ccgtgggtgac gcatgtcgcg caagactgta accacgcgtc 2040
 tgttgactgg cagtggtgag ccaatggtga tgtcagcgtt gaactgcgtg atgcggatca 2100
 acagtggttg gcaactggac aaggcactag cgggactttg caagtgggtg atccgcacct 2160
 ctggcaaccg ggtgaagggt atctctatga actgtgcgtc acagccaaaa gccagacaga 2220
 gtgtgatatc taccgccttc gcgtcggcat ccggtcagtg gcagtgaagg gccaacagtt 2280
 cctgattaac cacaaaccgt tctactttac tggcttttgt cgtcatgaag atgcggactt 2340
 acgtggcaaa ggattcgata acgtgctgat ggtgcacgac cacgcattaa tggactggat 2400
 tggggccaac tcctaccgta cctcgcatca cccttaacgt gaagagatgc tcgactgggc 2460
 agatgaacat ggcacgtggt tgattgatga aactgctgct gtccgcttta acctctcttt 2520
 aggcattggt ttccaagcgg gcaacaagcc gattaagag ctgatagcgc gtgacaaaaa 2580
 cggggaaact cagcaagcgc acttacaggg gattaaagag gatacccgtc cgcaagtgca 2640
 ccaccaagc gtggtgatgt gtagtattgc caacgaaccg gacccgacgc gtccgatcac 2700
 cgggaatatt tgcgcactgg cggaaagcaac gcgtaaactc gaccgcagcg gtccgatcac 2760
 ctgctcaat gtaatgttct gcgacgctca caccgatacc atcagcgatc tctttgatgt 2820
 gctgtgcctg aaccgttatt acggatggta tgtccaaagc ggogatttgg aaacggcaga 2880
 gaagggtact gaaaaagaac ttctggcctg gcaggagaaa ctgcatcagc cgattatcat 2940
 caccgaatac ggcgtggata cgttagccgg gctgcaactca atgtacaccg acatgtggag 3000
 tgaagagtat cagtgtgcat ggctggatat gtatcaccgc gtctttgatc gcgtcagcgc 3060
 cgtcgtcggg gaacaggtat ggaatttcgc cgattttgcg acctcgcaag gcatattgcg 3120
 cgttggcggg aacaagaaag ggatcttcac tcgcgacgcg aaaccgaagt cggcggcttt 3180

ccattcaggc	tgcgcaactg	ttgggaaggg	cgatcggtgc	gggcctcttc	gctattacgc	6840
cagctggcga	aagggggatg	tgctgcaagg	cgattaagtt	gggtaacgcc	aggggtttcc	6900
cagtcacgac	gttgtaaaac	gacggccagt	gccaagctga	cttggtcagc	ggccgcagat	6960
ttaggtgaca	ctata					6975

<210> 18
 <211> 3985
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:chimeric gene

<400> 18						
aagctttgct	ctagatcaaa	ctcacatcca	aacataacat	ggatatcttc	cttaccaatc	60
atactaatta	ttttggggtta	aataattaatc	attattttta	agatattaat	taagaaatta	120
aaagattttt	taaaaaaatg	tataaaatta	tattattcat	gatttttcat	acatttgatt	180
ttgataataa	atataatttt	tttaatttct	taaaaaatgt	tgcaagacac	ttattagaca	240
tagtcttggt	ctgttttaca	aagcattcat	catttaatac	attaaaaaat	atttaatact	300
aacagtagaa	tcttcttggtg	agtgggtgtg	gagtaggcaa	cctggcattg	aaacgagaga	360
aagagagtca	gaaccagaag	acaaataaaa	agtatgcaac	aaacaaatca	aaatcaaagg	420
gcaaaggctg	gggttggtctc	aattgggttg	tacattcaat	tttcaactca	gtcaacgggt	480
gagattcact	ctgacttccc	caatctaago	cgcggtatgca	aacggttgaa	tctaaccac	540
aatccaatct	cgttacttag	gggtttttcc	gtcattaact	cacctctgcc	accggtttc	600
cctataaaat	ggaactcaat	gctccccctc	aaactcgtat	cgcttcagag	ttgagacca	660
gacacactcg	ttcatatatac	tctctgctct	tctcttctct	tctacctctc	aaggtacttt	720
tcttctccct	ctaccaaatac	ctagattccg	tggttcaatt	tcggatcttg	cacttctggt	780
ttgctttgcc	ttgctttttc	ctcaactggg	tccatctagg	atccatgtga	aactctactc	840
tttctttaat	atctgcggaa	tacgcgttgg	actttcagat	ctagtcgaaa	tcatttcata	900
attgcctttc	tttcttttag	cttatgagaa	ataaaatcac	ttttttttta	tttcaaaata	960
aaccttgggc	cttggtgctga	ctgagatggg	gtttggtgat	tacagaattt	tagcgaattt	1020
tgtaattgta	cttggtttgtc	tgtagttttg	ttttgttttc	ttgtttctca	tacattcctt	1080
aggcttcaat	tttattcgag	tataggtcac	aataggaatt	caaactttga	gcaggggaat	1140
taatcccttc	cttcaaatcc	agtttggttg	tatatatgtt	taaaaaatga	aacttttgct	1200
ttaaattcta	ttataaacttt	ttttatggct	gaaatttttg	catgtgtctt	tgctctctgt	1260
tgtaaattta	ctgttttaggt	actaaactca	ggcttggtgt	gcagtttttg	aagtataacc	1320
atggtacgtc	ctgtagaaac	cccaaccogt	gaaatcaaaa	aactcgacgg	cctgtgggca	1380
ttcagctctg	atcgcgaaaa	ctgtggaatt	gatcagcggt	gggtgggaaag	cgcgttacaa	1440
gaaagccggg	caattgctgt	gccaggcagt	tttaacgata	agttcgccga	tgcatatatt	1500
cgtaattatg	cgggcaacgt	ctgggtatcag	cgcgaaagtct	ttataccgaa	aggttgggca	1560
ggccagcgta	tcgtgctgcg	tttcgatgcg	gtcactcatt	acggcaaaagt	gtgggtcaat	1620
aatcaggaag	tgatggagca	tcaggggcggc	tatacgccat	ttgaagccga	tgtcacgccc	1680
tatgttatgg	cggggaaaag	gttacgtatac	accgtttgtg	tgaacaacga	actgaactgg	1740
cagactatcc	cgcgggaat	gggtgattacc	gacgaaaacg	gcaagaaaaa	gcagtcttac	1800
ttccatgatt	tcttttaacta	tgccggaatc	catcgcagcg	taatgctcta	caccacgccc	1860
aacacctggg	tgagcgatat	caccgtgggtg	acgcattgtc	cgcaagactg	taaccacgcg	1920
tctgttgact	ggcaggtggg	ggccaatggg	gatgtcagcg	ttgaactgcg	tgatgcggat	1980
caacaggtgg	ttgcaactgg	acaaggcact	agcgggactt	tgcaagtggg	gaatccgcac	2040
ctctggcaac	cgggtgaagg	ttatctctat	gaactgtgcg	tcacagccaa	aagccagaca	2100
gagtggtgata	tctacccgct	tgcggtcggc	atccgggtcag	tggcagtga	gggccaacag	2160
ttcctgatta	accacaaacc	gttctacttt	actggctttg	gtcgtcatga	agatgcggac	2220
ttacgtggca	aaggattcga	taacgtgctg	atggtgcacg	accacgcatt	aatggactgg	2280
attggggcca	actcctaccg	tacctcgcat	tacctttacg	ctgaagagat	gctcgactgg	2340
gcagatgaac	atggcatcgt	gggtgattgat	gaaactgctg	ctgtcggctt	taacctctct	2400
ttaggcattg	gtttcgaagc	gggcaacaag	cggaaagaac	tgtacagcga	agaggcagtc	2460
aacggggaaa	ctcagcaagc	gcacttacag	gcgattaaag	agctgatagc	gcgtgacaaa	2520
aaccacccaa	gcgtggtgat	gtggagtatt	gccaacgaac	cggatacccg	tccgcaagtg	2580
cacgggaata	tttcgccact	ggcggaagca	acgcgtaaac	tcgaccgcac	gcgtccgac	2640
acctgcgtca	atgtaattgt	ctgcgacgct	cacaccgata	ccatcagcga	tctctttgat	2700
gtgctgtgcc	tgaaccgtta	ttacggatgg	tatgtccaaa	gcggcgattt	ggaaacggca	2760

gagaaggtag	tggaaaaaga	actttctggcc	tggcaggaga	aactgcatca	gccgattatc	2820
atcaccgaat	acggcgtgga	tacgttagcc	gggctgcact	caatgtacac	cgacatgtgg	2880
agtgaagagt	atcagtgtgc	atggctggat	atgtatcacc	gcgtctttga	tcgcgtcagc	2940
gccgtcgtcg	gtgaacaggt	atggaatttc	gccgattttg	cgacctcgca	aggcatattg	3000
cgcgttggcg	gtaacaagaa	agggatcttc	actcgcgacc	gcaaaccgaa	gtcggcggtc	3060
tttctgctgc	aaaaacgctg	gaactggcatg	aacttcggtg	aaaaaccgca	gcaggagggc	3120
aaacaatgaa	tcaacaactc	tcttggcgca	ccatcgctcg	ctacagcctc	ggtggggaat	3180
tccccggggg	tacctaatag	tgagatccaa	cacttacgtt	tgcaacgtcc	aagagcaa	3240
agaccacgna	cgcgggaagg	ttgcgcgagc	gtgtggattg	cgtctcaatt	ctctcttgca	3300
ggaatgcaat	gatgaatatg	atactgacta	tgaacttttg	aggggaatac	gcctagcacc	3360
gtcacctcat	aacgtgcata	atgcataccc	tgacaacatg	gaacatcgct	atctttctga	3420
agaattatgc	tcgttggagg	atgtcgcggc	aattgcagct	attgccaaca	tcgaactacc	3480
cctcacgcat	gcattcatca	atattattca	tgccgggaaa	ggcaagatta	atccaactgg	3540
caaatacatcc	agcgtgattg	gtaacttcag	ttccagcgac	ttgattcggt	ttgggtgctac	3600
ccacgttttc	aataaggacg	agatgggtga	gtaaaagaagg	agtgcgtcga	agcagatcgt	3660
tcaaacattt	ggcaataaag	tttcttaaga	ttgaatcctg	ttgccgggtc	tgcatgatt	3720
atcatataat	ttctgttgaa	ttacgttaag	catgtaataa	ttaacatgta	atgcatacgc	3780
ttatttatga	gatgggtttt	tatgattaga	gtcccgcaat	tatacattta	atacgcgata	3840
gaaaacaaaa	tatagcgcgc	aaactaggat	aaattatcgc	gcgcgggtgc	atctatgtta	3900
ctagatcgat	caaacttcgg	tactgtgtaa	tgacgatgag	caatcgagag	gctgactaac	3960
aaaaggtaca	tcggtcgcgc	agctc				3985

<210> 19

<211> 3684

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:chimeric gene

<400> 19

aagctttgct	ctagatcaaaa	ctcacatcca	aacataacat	ggatatcttc	cttaccaatc	60
atactaatta	ttttgggtta	aatattaatc	attattttta	agatattaat	taagaaatta	120
aaagattttt	taaaaaaatg	tataaaaatta	tattattcat	gattttttcat	acatttgatt	180
ttgataataa	atataatttt	tttaatttct	taaaaaatgt	tgcaagacac	ttattagaca	240
tagtcttgtt	ctgtttacaa	aagcattcat	catttaatac	attaaaaaat	atttaatact	300
aacagtagaa	tcttcttgtg	agtgggtgtg	gagtaggcaa	cctggcattg	aaacgagaga	360
aagagagtca	gaaccagaag	acaaataaaa	agtatgcaac	aaacaaatca	aaatcaaagg	420
gcaaaggctg	gggttggctc	aattgggttg	tacattcaat	tttcaactca	gtcaacgggt	480
gagattcact	ctgacttccc	caatctaagc	cgcggtatga	aacgggttga	tctaaccac	540
aatccaatct	cgttacttag	gggtttttcc	gtcattaact	cacccttgc	accgggttc	600
cctataaaat	ggaactcaat	gctcccctct	aaactcgtat	cgcttcagag	ttgagaccaa	660
gacacactcg	ttcatatctc	tctctgctct	tctcttctct	tctacctctc	aaggtaactt	720
tcttctccct	ctaccaaactc	ctagattccg	tggttcaatt	tcggatcttg	cacttctggt	780
ttgctttgcc	ttgctttttc	ctcaactggg	tccatctagg	atccatgtga	aactctactc	840
tttctttaat	atctgcggaa	tacgcgttgg	actttcagat	ctagtcgaaa	tcatttcata	900
attgcctttc	tttcttttag	cttatgagaa	ataaaatcac	ttttttttta	tttcaaaata	960
aaccttgggc	cttgtgctga	ctgagatggg	gtttgggtgat	tacagaattt	tagcgaattt	1020
tgtaattgta	cttgtttgtc	tgtagttttg	ttttgttttc	ttgtttctca	tacattcctt	1080
aggcttcaat	tttattcgag	tataggtcac	aataggaatt	caaactttga	gcaggggaat	1140
taatcccttc	cttcaaactc	agtttgtttg	tatatatgtt	taaaaaatga	aacttttgct	1200
ttaaattcta	ttataacttt	tttatggct	gaaatttttg	catgtgtctt	tgctctctgt	1260
tgtaaattta	ctgttttaggt	actaactcta	ggcttgttgt	gcagtttttg	aagtataacc	1320
atggccactt	tcttcgcccc	aacctccttc	ccctcccact	ctctctccaa	aaccttcgat	1380
acctatttcg	cccttgcccc	gaaagtcaac	gtctttgtga	acttcagggc	gaggaggcac	1440
gttgggggtg	gagtttcgaa	cgcgctgata	gaaccagatg	gagggaagct	cgtggagctt	1500
gtggtgacgg	attttgagag	ggatttgaag	aagggtgagg	ctctttcggt	gccgaggatc	1560
aagctctcaa	ggattgacct	tgagtgggtc	catgtcctca	gcgaaggatg	ggccacacc	1620
ctgaaaggct	tcatgagaga	agccgagttc	ctccaaacgc	ttcatttcaa	ctcgctccga	1680
ctcgatgatg	ggtcggctgt	gaacatgtca	gtgcccacgc	tgctggctat	tgatgatgcg	1740

cagaagcatc	ggatcgggga	taacaaaaag	gttgctcttt	ttgattccaa	gggagacccc	1800
gttgcaattc	tcaataatat	tgagatttat	aagcatccta	aagaagaaag	aatagcccga	1860
acttggggaa	ccattgcccc	tggcctacct	tatgttgaa	aaactataac	caatgctgga	1920
aattggttga	ttgggggtga	cctagagggtc	attgaaccaa	ttcagtacaa	tgatggactt	1980
gatcattttc	gtctatctcc	ggcacaactc	cgtgcagagt	tcacaaggcg	caatgcggat	2040
gctgtgtttg	ccttccagct	ccggaatcct	gttcacaatg	gccatgcttt	gctaatagact	2100
gacacccgaa	agcgccttct	tgagatgggc	tataagaatc	ctgtcctctt	gcttcatcca	2160
cttgagggtc	acaccaaagc	tgatgatgtc	ccacttgatt	ggcgaatgaa	gcaacatgag	2220
aagggtactg	aggatgggtg	tcttgatcca	gagacaactg	tggtatccat	attcccatct	2280
cccatgcact	atgctggacc	cacggagggtg	cagtggcatg	caaaggctag	gatcaatgca	2340
ggggctaact	tctatatcgt	tggtcgtgac	cccgcaggca	tgagccatcc	agttgagaaa	2400
agagatctgt	atgatgctga	ccatggaaag	aaagtattga	gcatggcacc	gggactagag	2460
cgtctaaaca	ttcttccctt	caggggttgc	gcatatgaca	agactcaggg	taaaatggca	2520
ttctttgacc	cttcaaggcc	tcaggacttc	ctgttcata	caggcacaaa	gatgcgcaca	2580
ctggcaagga	acaaagaaag	tcttccctgat	ggatttatgt	gccctgggtg	atggaagggtg	2640
ctggttgatt	actatgatag	cttagtactc	tcaagcaacg	gcaaagtgca	ggaagctgtt	2700
ccagcttaat	cttgtatcat	atcataatgt	atatatctca	tgattgggag	aaaccttaag	2760
cttatgtatt	ctcctgctaa	gacataactc	acgaggatcc	tctggcccaa	tctaataata	2820
ataataaatt	aaaacttttg	ggaggcaaaa	aaaaaaaaaa	aaaaaaaaaa	aactcgaggg	2880
ggggcccggt	acctaatagt	gagatccaac	acttacgttt	gcaacgtcca	agagcaaata	2940
gaccacgnac	gccggaaggt	tgccgcagcg	tgtggattgc	gtctcaattc	tctcttgacg	3000
gaatgcaatg	atgaatatga	tactgactat	gaaactttga	gggaatactg	cctagcaccg	3060
tcacctcata	acgtgcatca	tgcatgccct	gacaacatgg	aacatcgcta	tttttctgaa	3120
gaattatgct	cgttgaggga	tgctgcggca	attgcagcta	ttgccaacat	cgaactaccc	3180
ctcacgcgat	cattcatcaa	tattattcat	gcggggaaag	gcaagattaa	tccaactggc	3240
aaatcatcca	gcgtgattgg	taacttcagt	tccagcgact	tgattcgttt	tggtgctacc	3300
cacgttttca	ataaggacga	gatgggtggag	taaagaagga	gtgcgtcgaa	gcagatcggt	3360
caaacatttg	gcaataaagt	ttcttaagat	tgaatcctgt	tgccggtctt	gcgatgatta	3420
tcatataaatt	tctgttgaat	tacgttaagc	atgtaataat	taacatgtaa	tgcatgacgt	3480
tatttatgag	atgggttttt	atgattagag	tcccgcatt	atacatttaa	tacgcgatag	3540
aaaacaaaat	atagcgcgca	aactaggata	aattatcgcg	cgcggtgtca	tctatgttac	3600
tagatcgatc	aaacttcggt	actgtgtaat	gacgatgagc	aatcgagagg	ctgactaaca	3660
aaaggtacat	cggtcgacga	gctc				3684

<210> 20

<211> 3963

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:chimeric gene

<400> 20

aagctttgct	ctagatcaaa	ctcacatcca	aacataacat	ggatatcttc	cttaccaatc	60
atactaatta	ttttgggtta	aatattaatc	attattttta	agatattaat	taagaaatta	120
aaagattttt	taaaaaaatg	tataaaatta	tattattcat	gatttttcat	acatttgatt	180
ttgataataa	atataatttt	tttaatttct	taaaaaatgt	tgcaagacac	ttattagaca	240
tagtcttggt	ctgtttacaa	aagcattcat	catttaatac	attaaaaaat	atttaatact	300
aacagtagaa	tcttcttggtg	agtgggtgtg	gagtaggcaa	cctggcattg	aaacgagaga	360
aagagagtca	gaaccagaag	acaaataaaa	agtatgcaac	aaacaaatca	aaatcaaagg	420
gcaaaggctg	gggttggtgc	aattgggtgc	tacattcaat	tttcaactca	gtcaacggtt	480
gagattcact	ctgacttccc	caatctaagc	cgcggtatgca	aacggttgaa	tctaaccac	540
aatccaatct	cgttacttag	gggtttttcc	gtcattaact	cacctctgcc	acccggtttc	600
cctataaatt	ggaactcaat	gtccttctct	aaactcgtat	cgcttcagag	ttgagaccaa	660
gacacactcg	ttcatatata	tctctgctct	tctcttctct	tctacctctc	aaggtaactt	720
tcttctccct	ctaccaaata	ctagattccg	tggttcaatt	tcggatcttg	cacttctggt	780
ttgctttgcc	ttgctttttc	ctcaactggg	tccattcagg	atccatgtga	aactctactc	840
tttctttaat	atctgcggaa	tacgcgttgg	actttcagat	ctagtcgaaa	tcatttcata	900
attgcctttc	tttcttttag	cttatgagaa	ataaaatcac	ttttttttta	tttcaaaata	960
aaccttgggc	cttgtgctga	ctgagatggg	gttttggtgat	tacagaattt	tagcgaattt	1020

tgtaattgta	cttgtttgtc	tgtagttttg	ttttgttttc	ttgttttctc	tacattcctt	1080
aggcttcaat	tttattcgag	tataggtoac	aataggaatt	caaactttga	gcagggggaat	1140
taatcccttc	cttcaaattc	agtttgtttg	tatatatggt	taaaaaatga	aactttttgct	1200
ttaaaattcta	ttataaacttt	ttttatgggt	gaaatttttg	catgtgtctt	tgctctctgt	1260
tgtaaaattta	ctgttttaggt	actaactcta	ggcttggtgt	gcagtttttg	aagtataacc	1320
atggccggtt	cgagctcgca	catgcgtttc	acctttgagt	gccgctccga	tcccgatttc	1380
tcgccccccc	cgccgtcctt	cgacaacctc	cgccgcgcaa	acttccgctc	ctccgcagga	1440
tccgggcgcg	cgtttcacgg	catctcctcc	ctcatcctcc	gcttccctcc	caacttccag	1500
cgccagctaa	gcaccaaggc	gcgcgcgcaac	tgagcaaca	tcggcgctcg	gcaaactcgtc	1560
gccgcttcgt	ggtcgaacaa	cagcgacaac	tctccggccg	ccggggctcc	ggcgccgccc	1620
gcggccaccg	ccacggacgc	cgtacgggtg	cctctccccg	tcgtcgctcg	cgccaacgag	1680
gacgtcgttg	tctccgccgc	ggcagacgag	aacggggctg	tacagttaaa	cagtgttct	1740
tattcttcat	ttttgaaatc	cgatgcaagc	aaaacgattc	atgccgctga	aagactgggt	1800
aggggtattg	agactgatgg	aattaccacc	cctgtgggtt	acacttctgc	ctactttttt	1860
aagaaaaccg	ctgatctcat	tgatttcaag	gagaatcgtc	aagtgaagta	tgaatacggg	1920
cgctatggaa	acccaacgac	ggtggttctg	gaggagaaga	taagtgcatt	ggagggggcc	1980
gaatcaactg	tgataatggc	gtctgggatg	tgtgctagcg	tagtcctgtt	tatggcactg	2040
gttccagctg	gtggacatct	tgtgaccact	acggattgtt	ataggaagac	tagaatattc	2100
attgagactt	ttcttccaaa	gatggggatc	acgaccactg	taattgatcc	agcagatgtt	2160
ggagccttgg	aatctgcatt	ggagcagcac	aatgtgtctc	tattcttcac	tgagtctcct	2220
accaatccat	tcctgagatg	tgttgatatt	aagctgggtt	cagagctttg	ccacaagaag	2280
gggactttgc	tctgtattga	tggtagattt	gcaactccat	tgaaccagaa	ggcccttgcc	2340
cttggcgctg	atctgattct	gcactcctta	acaaaataca	tgggtggaca	tcatgatgtc	2400
cttgggtggt	gcataagtgg	ttcaattaag	gtgggttctc	aaattcggac	tttgacccat	2460
gttttggttg	gtacacttaa	ccgaatgct	gcatacctat	tcatacagag	catgaaaacg	2520
ctgcatctcc	gtgtacagca	gcagaattca	acaggaatga	ggatggccaa	acttttagag	2580
gcacatccca	aggtgaagcg	ggtctactat	ccaggcttgc	cgagtcaccc	tgaacatgag	2640
cttgccaaga	ggcagatgac	tggtttcggt	ggtgttgtca	gttttgagat	tgatggagat	2700
ctacatacca	caataaaaatt	tattgattca	ttgaaaatcc	catatattgc	ggcctcggtt	2760
ggtggctgtg	agagcattgt	ggatcaacct	gctattttgt	cttactggga	tcttctcag	2820
tcagaaaggg	ccaagtacaa	gatttatgac	aacctgggtc	gcttcagctt	tggagttgaa	2880
gattttgagg	atttgaaggc	tgatgtcctg	caagctctgg	aagctatata	gacagttttc	2940
ctgattcacc	caagtttttt	tcttttataa	ttgtgctatt	tgtttggtat	cacatctggc	3000
gattcaattg	aattttgatc	gtctaattgt	ctgttggaat	tgtgttaaga	tgaatggctc	3060
ctaatttgga	tgttatgaaa	cttgtgatga	attgttgaaa	ttgaaacctc	tatttgatga	3120
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	actcgagggg	gggcccggta	cctaatagtg	3180
agatccaaca	cttacgtttg	caacgtccaa	gagcaaatag	accacgnacg	ccggaagggt	3240
gccgcagcgt	gtggattgcg	tctcaattct	ctcttgacag	aatgcaatga	tgaatatgat	3300
actgactatg	aaactttgag	ggaatactgc	ctagcaccgt	cacctcataa	cgtgcatcat	3360
gcatgccctg	acaacatgga	acatcgctat	ttttctgaag	aattatgctc	gttgagggat	3420
gtcgcggcaa	ttgcagctat	tgccaacatc	gaactacccc	tcacgcagtc	attcatcaat	3480
attattcatg	cggggaaagg	caagattaat	ccaactggca	aatcatccag	cgtgattggg	3540
aacttcagtt	ccagcgactt	gattcgtttt	ggtgctaccc	acgttttcaa	taaggacgag	3600
atggtggagt	aaagaaggag	tgcgtcgaag	cgatcgcttc	aaacatttgg	caataaagtt	3660
tcttaagatt	gaatcctggt	gccggtcctg	cgatgattat	catataattt	ctggtgaatt	3720
acgttaagca	tgtaataaatt	aacatgtaat	gcatgacgtt	atztatgaga	tgggttttta	3780
tgattagagt	cccgaatta	tacatttaatt	acgcgataga	aaacaaaata	tagcgcgcaa	3840
actaggataa	attatcgcg	gcggtgtcat	ctatgttact	agatcgatca	aacttcggta	3900
ctgtgtaatg	acgatgagca	atcgagaggg	tgactaacaa	aaggtacatc	ggtcgacgag	3960
ctc						3963